



# Efficiency of N<sub>2</sub> Gas Flushing Compared to the Lactoperoxidase System at Controlling Bacterial Growth in Bovine Raw Milk Stored at Mild Temperatures

Patricia Munsch-Alatossava<sup>1\*</sup>, Romanie Quintyn<sup>1</sup>, Ingrid De Man<sup>2</sup>, Tapani Alatossava<sup>1</sup> and Jean-Pierre Gauchi<sup>3</sup>

<sup>1</sup> Department of Food and Environmental Sciences, University of Helsinki, Helsinki, Finland, <sup>2</sup> Food Technology, Vives University College Campus Roeselare, Roeselare, Belgium, <sup>3</sup> MalAGE, INRA, Université Paris-Saclay, Jouy-en-Josas, France

## OPEN ACCESS

### Edited by:

Javier Carballo,  
University of Vigo, Spain

### Reviewed by:

Carmen Wacher,  
Universidad Nacional Autónoma de  
México, Mexico  
Robert Lawrence Brown,  
United States Department of  
Agriculture, Agricultural Research  
Service, USA

### \*Correspondence:

Patricia Munsch-Alatossava  
patricia.munsch@helsinki.fi

### Specialty section:

This article was submitted to  
Food Microbiology,  
a section of the journal  
Frontiers in Microbiology

**Received:** 29 March 2016

**Accepted:** 19 May 2016

**Published:** 02 June 2016

### Citation:

Munsch-Alatossava P, Quintyn R, De Man I, Alatossava T and Gauchi J-P (2016) Efficiency of N<sub>2</sub> Gas Flushing Compared to the Lactoperoxidase System at Controlling Bacterial Growth in Bovine Raw Milk Stored at Mild Temperatures. *Front. Microbiol.* 7:839. doi: 10.3389/fmicb.2016.00839

To prevent excessive bacterial growth in raw milk, the FAO recommends two options: either cold storage or activation of the lactoperoxidase system (LPs/HT) in milk with the addition of two chemical preservatives, hydrogen peroxide (H) and thiocyanate (T). N<sub>2</sub> gas flushing of raw milk has shown great potential to control bacterial growth in a temperature range of 6–12°C without promoting undesired side effects. Here, the effect of N<sub>2</sub> gas (N) was tested as a single treatment and in combination with the lactoperoxidase system (NHT) on seven raw milk samples stored at 15 or 25°C. For the ratio defined as bacterial counts from a certain treatment/counts on the corresponding control, a classical Analyse of Variance (ANOVA) was performed, followed by mean comparison with the Ryan-Einot-Gabriel-Welsch multiple range test (REGWQ). Altogether, the growth inhibition was slightly but significantly higher at 25°C than at 15°C. Except for one sample, all ratios were lower for HT than for N alone; however, these differences were not judged to be significant for five samples by the REGWQ test; in the remaining two samples, N was more effective than HT in one case and less effective in the other case. This study shows that N<sub>2</sub> gas flushing, which inhibited bacterial growth in raw milk at 15 and 25°C for 24 and 12 h, respectively, could constitute an alternative to LPs where no cold storage facilities exist, especially as a replacement for adulterating substances.

**Keywords:** raw milk, N<sub>2</sub> gas, lactoperoxidase system, antibacterial, Ryan-Einot-Gabriel-Welsch test

## INTRODUCTION

Despite high-tech food production systems, in so-called developed countries, approximately 30–40% of food is lost during retail, in food service or at home; the losses equal those from developing countries, which are mostly attributable to the absence of food chain infrastructure (Gustavsson et al., 2011). In Europe, as much as 20% of the dairy products are lost or wasted every year (FAO, 2012). The presence of spoilage microorganisms, antibiotic-resistant bacteria and the risk of food-borne pathogens are a great challenge to food production systems. Although, milk is considered to be sterile when secreted from a healthy udder, numerous contamination sources increase its bacterial load; hence, depending on the production area, the farm type and the milk



(Munsch-Alatossava et al., 2010a). All raw milk samples (100 ml per bottle and treatment) were continuously mixed and stored at 15 or 25°C ± 0.1 in a water bath until subjected to analyses. The description of the conditions (applied to every sample) is as follows: C = non treated milk (accounts for the control); T = SCN<sup>-</sup>; H = H<sub>2</sub>O<sub>2</sub>; HT = H<sub>2</sub>O<sub>2</sub> + SCN<sup>-</sup>; N = N<sub>2</sub> flushing; NT = N<sub>2</sub> flushing + SCN<sup>-</sup>; NH = N<sub>2</sub> flushing + H<sub>2</sub>O<sub>2</sub>; NHT = N<sub>2</sub> flushing + H<sub>2</sub>O<sub>2</sub> + SCN<sup>-</sup>.

## Microbiological Analyses

For the analyses, at the given sampling points, the raw milk was serially diluted in 0.85% NaCl solution. “Total bacterial counts,” Gram-negative bacteria, lactobacilli and streptococci/lactococci were determined mostly from triplicate (if not duplicate) platings on Plate Count Agar (PCA, LabM, Ltd, Lancashire, UK), MacConkey agar (MC, Becton Dickinson, Le Pont de Claix, France), MRS agar (LabM, Ltd, Lancashire, UK), and M17 agar (Sigma-Aldrich, Steinheim, Germany), respectively, after incubation at 30°C for 3 days under aerobic conditions for PCA and MC plates or at 37°C for 72 h for MRS and M17 plates under anaerobiosis. Anaerobic culture jars and AnaeroGen<sup>TM</sup> sachets (Oxoid Ltd, Basingstoke, UK) were used for creating the anaerobic incubation conditions.

## Statistical Analyses

To compare the efficiency of the different treatments, the analyses considered the ratio defined as “the counts obtained on a certain treatment divided by the counts enumerated on the corresponding control.” The statistical analyses determined the significance of the effects of the three considered factors (“time”/duration of the treatment, “media type” and “condition”/applied treatment); moreover, two-factor and three-factor interactions on the ratios were evaluated. At first, a classical ANOVA (Analysis of Variance) was performed. Considering the *p*-values associated with the effects and interactions (*p* < 0.001) and because the effect of the factor “condition” was always found to be highly significant, a comparison of the ratio means that reflect the response to the seven treatments (T, H, HT, N, NT, NH, and NHT) was then undertaken with the Ryan-Einot-Gabriel-Welsch test (REGWQ) for multiple comparison of means, as described by Hsu (1996), with an alpha risk of 0.05. All calculations were performed with the SAS/STAT software version 9.4/ GLM procedure (SAS Institute, NC, USA).

## RESULTS

### Ranking of the Treatments Comparison of Raw Milk Samples

The ANOVA revealed a strong significant “sample effect” indicating that the mean ratios in response, to the seven treatments were significantly different (data not shown). Subsequently, the Ryan-Einot-Gabriel-Welsch (REGWQ) multiple range test, applied to the ratios, led to a significant grouping of the seven raw milk samples (M1–M7) into three categories, depending on the combined inhibitory effects caused by the LPs- and N<sub>2</sub>-based treatments (Table 1). The treatments induced the most contrasting effects for the three samples

**TABLE 1 | REGWQ ranking of the seven bovine raw milk samples stored at 15 and 25°C and treated with LPs and N<sub>2</sub> gas.**

Raw milk sample/Applied storage temperature/Start of the analysis	Mean ratio <sup>a</sup>	Significance <sup>b</sup>
M1/15°C/Apr 13, 2015	0.7795	A
M3/15°C/Jun 10, 2015	0.5728	B
M6/25°C/May 25, 2015	0.5324	B
M7/25°C/Jun 8, 2015	0.5030	B
M5/25°C/May 4, 2015	0.4251	B C
M4/25°C/Apr 27, 2015	0.2812	C
M2/15°C/Apr 20, 2015	0.2518	C

<sup>a</sup>Mean of the ratios from the seven different examined conditions. <sup>b</sup>Means with the same letter are not significantly different (alpha risk = 0.05).

considered in April as the inhibition of bacterial growth was highest in samples M2 and M4 and lowest for M1.

The four samples stored at 25°C followed each other in the ranking. The efficiency of the treatments to trigger bacterial inhibition appeared to be more dependent on the sample than on the applied temperature as both the highest and lowest inhibitory effects were recorded for two samples (M1 and M2) stored at 15°C (Table 1). The treatments showed a rather equivalent inhibitory effect, -with a difference of 0.0294- for the tightly ranked samples M4 and M2, although they were stored at 25 and 15°C, respectively, which equaled the difference recorded for samples M6 and M7, both as tightly ranked and stored at 25°C (Table 1). Altogether, the treatments proved to be slightly but significantly more inhibitory at 25°C than at 15°C.

### Raw Milk Storage at 15°C

The REGWQ test distinguished four categories among the tested conditions depending on their impact on the three milk samples (M1, M2, M3) that were stored at 15°C (Table 2). The sample M2, which presented the highest initial “total counts” (4 log-units) and for which the ratio “counts on MacConkey agar/total counts” was lowest, showed the highest sensitivity to the treatments as the ratios were below 0.15 for five conditions; in contrast, sample M3 presented the lowest initial counts (2.6 log-units) but less inhibition was observed based on the ratios (Table 2). Based on counts from all four agar medium types, condition T was ranked apart in all three experiments with two ratios exceeding 1, indicating that the addition of SCN<sup>-</sup> exerted a clear stimulatory effect on bacterial growth for samples M1 and M3 (Table 2). The addition of H<sub>2</sub>O<sub>2</sub> alone (H) triggered an inhibition of approximately 30% for M1 and M3 and of 62% for M2 (Table 2). The activated lactoperoxidase system (HT) induced variable levels of inhibition: 18, 95, and 67% for M1, M2, and M3, respectively (Table 2). For N<sub>2</sub> flushing (N), the recorded inhibitory percentages were 62, 86, and 38% for M1, M2, and M3, respectively. The difference between N and HT was significant for M1, where N alone supplanted HT; but HT and N were equally efficient for M2 and M3. In all three experiments, the ratio was significantly lower for NT than for T; it was different from N for M1, and similar for M2 and M3, suggesting that the inhibitory effects of N<sub>2</sub> gas flushing overcame the stimulatory effect of

**TABLE 2 | Results of the Ryan-Einot-Gabriel-Welsch multiple range test for the three raw milk samples stored at 15°C.**

Raw milk sample	Initial “total counts” <sup>a</sup>	Initial counts on MacConkey agar/“total counts”	Condition	Ratio	Significance <sup>b</sup>	F-value of the most important factors
M1	3.9	ND <sup>c</sup>	T	1.7033	A	Time 32.44
			NT	1.0329	B	Condition 31.02
			HT	0.8197	B C	Medium 27.12
			H	0.7029	C D	
			NHT	0.4540	D	
			N	0.3817	D	
			NH	0.3650	D	
M2	4.0	0.9	T	0.9243	A	Condition 112.23
			H	0.3832	B	Medium 35.15
			NT	0.1437	C	
			N	0.1390	C D	
			NH	0.0547	C D	
			HT	0.0488	C D	
			NHT	0.0127	D	
M3	2.6	1.1	T	1.2316	A	Time 28.02
			H	0.6821	B	Condition 16.84
			N	0.6243	B C	Medium 11.75
			NH	0.5148	B C	
			NT	0.4590	B C D	
			HT	0.3337	C D	
			NHT	0.1716	D	

<sup>a</sup>“Total counts” on PCA, <sup>b</sup>Mean ratios with the same letter are not significantly different (alpha risk = 0.05), <sup>c</sup>ND = not determined.

T (Table 2); for all three experiments, the effects due to the combination of N and H (NH) did not differ significantly from those observed for N alone. Compared to H alone, the addition of N to H increased the inhibitory effect for M2 (Table 2). For both milk samples M2 and M3, the inhibitory effect was maximal for condition NHT but not significantly different from condition HT (for M2 and M3) or from N (for M2).

### Raw Milk Storage at 25°C

The REGWQ test revealed that the treatments were less discriminatory at 25°C, as only three categories were distinguished among the applied treatments for the four milk samples. However, despite greater homogeneity in initial “total counts” (which ranged between 3.1 and 3.8 log-units), the four samples responded differently to the treatments (Table 3).

For raw milk samples stored at 25°C, T also stimulated bacterial growth as the mean ratio was highest for M4 and it exceeded 1 for M5, M6, and M7 (Table 3). Condition H yielded ratios undistinguishable from T for M5 and M6, but it showed a moderate inhibitory effect for M4 and M7 (Table 3). HT promoted the highest growth inhibition for samples M6 and M7. If higher inhibitory ratios were recorded for HT compared to N<sub>2</sub> flushing (N), these differences were not judged to be significant for three samples. At 25°C, with the exception of M7, N<sub>2</sub> flushing alone (N) achieved a reduction in bacterial levels similar to that obtained with HT (Table 3). NHT promoted the highest inhibition of bacterial growth for samples M4 and M5.

Altogether, NHT was tightly ranked with HT as the differences between the two conditions were not significant for any of the investigated raw milk samples (Table 3).


### Contribution of the Major Factors That Determined the Ranking

The analysis of the contribution of the different factors to the ranking of the ratios revealed that the “condition” (or applied treatment), the storage “time” of the milk and the “medium type” affected the seven raw milk samples (M1–M7) differently (Tables 2, 3). Altogether, although the factors varied in their respective importance, the “condition” was a major determinant for four samples (M2, M4, M5, and M6), with an overwhelming impact on the ranking of the treatments for sample M2. Clearly, the factor storage “time” also displayed great heterogeneity among the seven raw milk samples as it mainly determined the ranking of the treatments for samples M3 and M7; altogether, its effect was variable and more moderate as noticeable for M1, for example, where the three factors played an equivalent role (Tables 2, 3). Considering the factor storage “time” during the course of the experiments, at 15°C the inhibitory effects increased during storage for M1, they were high and constant for M2, and they fluctuated over time for M3 being maximal at 24 and 48 h storage (Figure 1A). At 25°C, for samples M5 and M6, the inhibition was maximal at 12 h, then the mean ratios increased over time, indicating declining inhibition; for M7, the highest

**TABLE 3 | Results of the Ryan-Einot-Gabriel-Welsch multiple range test for the four raw milk samples stored at 25°C.**

Raw milk sample	Initial “total counts” <sup>a</sup>	Initial counts on MacConkey agar/“total counts”	Condition	Ratio	Significance <sup>b</sup>	F-value of the most important factors
M4	3.6	ND <sup>c</sup>	T	0.9027	A	Condition 40.30 Medium 9.19
			H	0.6019	B	
			N	0.1823	B C	
			NT	0.1036	C	
			NH	0.0572	C	
			HT	0.0521	C	
			NHT	0.0398	C	
M5	3.2	0.8	H	1.0630	A	Condition 46.01 Time 27.87 Medium 7.74
			T	0.8939	A B	
			NT	0.6955	B	
			N	0.1733	C	
			NH	0.1060	C	
			HT	0.0271	C	
			NHT	0.0130	C	
M6	3.8	0.9	T	1.2494	A	Condition 23.93 Time 14.33
			H	1.1004	A	
			NH	0.3955	B	
			NT	0.3750	B	
			N	0.3324	B	
			NHT	0.1311	B	
			HT	0.1185	B	
M7	3.1	1.2	T	1.2041	A	Time 26.19 Condition 22.64 Medium 12.56
			H	0.5957	B	
			N	0.5279	B	
			NH	0.4186	B C	
			NT	0.3541	B C	
			NHT	0.2172	C	
			HT	0.1954	C	

<sup>a</sup>Total counts” on PCA, <sup>b</sup>Mean ratios with the same letter are not significantly different (alpha risk = 0.05), <sup>c</sup>ND = not determined.



**FIGURE 1 | Impact of the factor “storage time in h” on the ratios determined for raw milk stored at 15°C (A) (for 12–72 h), or at 25°C (B) (for 12–24 h).**

inhibition occurred at intermediate storage time (17 h), whereas for M4 the inhibition remained unchanged for 24 h. (**Figure 1B**).

Consideration of the factor “medium type” (which supported the growth of different bacterial populations, together with the corresponding incubation conditions) showed that altogether the different bacterial population types were not similarly affected


by the treatments at either considered storage temperature (**Figure 2**). For M1, bacterial types on MRS and M17 agar were recovered in high amounts under the treatments compared to PCA and MC whereas for M2, clearly all population types were hindered but mainly those enumerated on PCA and on MC, in contrast to M3 where populations on MRS were mostly

inhibited (**Figure 2A**). At 25°C, the inhibitory effects recorded on the different agar types also showed milk-sample dependent responses. “Total counts” on PCA were inhibited by 85% for M4 but only by 35% for M6. Gram-negative bacterial types were also mostly inhibited for M4 (75%), whereas the inhibition ranged from 60% (M6 and M7) to 45% for M5. The inhibition of lactobacilli varied greatly between 70% (M4) and 25% (M7), in contrast to streptococci/lactococci for which the inhibitory effects (which ranged between 42 and 52%) were mostly similar between all four samples (**Figure 2B**).


ANOVA tables revealed double and triple interactions between the considered factors storage “time,” “medium” type, and “condition,” which were often significant (at an alpha risk factor of 5%) but generally lower than the main effects due to the three factors alone (data not shown).

## Trends of Bacterial Growth for Samples M2 and M5

The detailed results of the bacterial growth trends for samples M2 and M5, which exhibited high sensitivity to the treatments (Category C in **Table 1**), are shown in **Figures 3, 4**. For sample M2 (stored at 15°C), irrespective of the sampling time points, similar bacterial levels were enumerated on PCA for condition T to those of the control C (**Figure 3A**). Compared to C, at 24 h, inhibition of growth was observed for H (−1.8 log-units), for N (−2.0 log-units), for NT (−1.9 log-units) and for NH (−2.1 log units); conditions HT and NHT yielded lower bacterial levels (−2.8 and −2.7 log-units, respectively) than C. At the 48 and 72 h time points, the “total bacterial counts” for T and H were quite similar to those of C in contrast to all the other conditions (HT, N, NT, NH, and NHT), which showed 1–1.5 log-units less bacteria




**FIGURE 2 |** Ratio levels illustrating the incidence of the treatments on bacterial population types enumerated on four considered agar types for raw milk stored at 15°C (A) or at 25°C (B). PCA, (Plate Count Agar); MC, (MacConkey agar); MRS, (culture of lactic acid bacteria); M17, (lactic streptococci).



**FIGURE 3 |** Bacterial counts determined for sample M2 stored at 15°C on (A) PCA (Plate Count Agar), (B) MacConkey agar, (C) MRS agar, (D) M17 agar. Error bars indicate standard deviations based mainly on three replicates.





after 48 h; however, at 72 h, the inhibitory effects faded and only condition NHT showed lower counts (Figure 3A). Of note, like HT, N kept the counts below 5.5 log-units for more than 24 h. For M1 and M3, a similar level of inhibition was observed in one case, and increased inhibition was observed for the other case (data not shown). Over time, on MC agar, C and T showed similar bacterial growth trends (Figure 3B). In contrast, H exhibited counts close to 2 log-units lower than C at 24 h; but the final counts for H were only slightly lower at 48 and 72 h than for C or T. A bactericidal type of action was observed for HT as counts dropped during the first 24 h then increased thereafter. All conditions where milk samples were N<sub>2</sub>-flushed showed that bacterial counts were below 5.5 log-units after 24 h of storage; this observation was still valid at 48 h for NH and NHT, but counts slightly increased for N (5.6 log-units) and NT (5.8 log-units). Notably at 48 and 72 h, the counts were slightly lower for NHT than for HT (Figure 3B). The hindrance of the growth of lactobacilli was only moderate as all conditions showed only slightly lower bacterial levels than C or T at 24 h (Figure 3C). Compared to HT and N, NHT exhibited lower counts after 48 and 72 h. For streptococci/lactococci, all treatments showed approximately 1 log-unit lower counts at 24 h on M17, except for T, which was indistinguishable from C (Figure 3D). At 48 h, all counts had increased similarly except for HT and NHT for which some delay in growth was observed. At 72 h, the counts were approximately 1.8 log-units lower for NHT than for C (Figure 3D).

For M5, after 12 h storage at 25°C, conditions C, H, and T exhibited over 6.5 log-units “total counts” on PCA, whereas all

other conditions kept bacterial levels below 5.5 log-units. At 17 h, counts from HT, NH, and NHT were below or approximately at 5.5 log-units, whereas they reached 5.9 log-units for N (Figure 4A). A slight inhibitory effect was still noticeable at 24 h for conditions HT and NHT. On MC, after 12 h storage at 25°C, Gram-negative bacterial counts had increased greatly for C and T; H and N alone or in combination with T (NT) yielded counts that ranged between 4.7 and 5.2 log-units; at 24 h, HT and NHT showed lower bacterial counts (Figure 4B). On MRS, inhibited lactobacilli growth was mainly observed with HT and NHT as counts were lowest at 12, 17, and 24 h (Figure 4C). If HT, in contrast to N inhibited the growth of lactobacilli during the initial 12 h of storage, at 17 h, the level was equivalent for HT and N. On M17, streptococcal/lactococcal counts increased similarly for C, T, H, N, and NT (Figure 4D), in contrast to HT, NH and NHT where counts were lower after 12 h of storage. At 17 h, some inhibitory effect was still noticed for HT, N, NH, and NHT, as it was for HT and NHT at 24 h (Figure 4D).

## DISCUSSION

In this study, N<sub>2</sub> gas flushing and chemical preservatives (SCN<sup>-</sup> and H<sub>2</sub>O<sub>2</sub> which serve to activate the lactoperoxidase system (LPs) in raw milk) were used to treat seven lorry raw milk samples (M1–M7) that presented a “certain” bacterial load and diversity resultant from a “certain” cold storage history; as “total counts” ranged between 2.6 and 4.0 log-units at the initial stage, all seven raw milk samples can be considered of very good bacteriological

quality (Tables 2, 3) as “total bacterial counts” per ml in the tank at the farm and at the dairy silo level should not exceed  $10^5$  and  $3.10^5$  cfu/ml (5.48 log-units), respectively (Anonymous, 2004).

The treatments were limited to 24 h at 25°C and to 72 h at 15°C when also most treated milk samples exceeded 5.5 log-units in “total counts” (Figures 3A, 4A and data not shown). Notably, despite a 10°C difference in the applied storage temperature, a certain similarity between the inhibitory patterns was noticeable for M2 and M4, and for M3 and M6 (with the exception of lactobacilli, not determined) (Figure 2 and data not shown), suggesting that populations could be similarly targeted by the treatments irrespective of the storage temperature.

Altogether, the treatments were most efficient for samples M2, M4, and M5 (Table 1). Analysis of the factors that impacted the ranking of the raw milk samples revealed that for these samples, the factor “condition” was preponderant in the ranking (Tables 2, 3).

Considering the values of the ratios that reflect the inhibition due to the treatments, altogether, the inhibitory effects of either the LPs or the N<sub>2</sub> gas flushing were more dependent on the sample (or initial microbiota) than the initial bacterial level (Tables 1, 3); this point is well illustrated by the position of sample M2 in the ranking (Table 1) which showed the highest sensitivity to the applied treatments despite the fact that it presented the highest initial bacterial counts. Perhaps this result could be explained by better synergy with the indigenous LPs in this raw milk sample or by the presence of other bacterial inhibitors (Korhonen and Marnila, 2002; Marnila and Korhonen, 2002).

The microbiological analyses showed that bacterial counts on MacConkey agar/“total counts” on PCA agar were lowest for M2, M5, and M6, and higher for M3 and M7 (Tables 2, 3). Determination of the inhibitory ratios revealed that for treatments HT and N, the values were lowest for samples M2, M5, and M6 and highest for M3 and M7 (Tables 2, 3). It is well known that cold storage of raw milk promotes an increase of Gram-negative bacterial types (Chambers, 2002); altogether, our data suggest that both HT and N treatments achieved the highest inhibition of bacterial growth in milk samples not yet totally dominated by Gram-negative bacteria.

The performance of the activated lactoperoxidase system (HT) enabled bacterial growth to be controlled for 24 h in raw milk stored at 15°C (for M1 until 48 h) (Figure 3A and data not shown), and for 12 h in milk stored at 25°C (Figure 4A and data not shown), respectively, which is in line with previous reports (FAO, 2005). When the milk was stored at 15°C, the N<sub>2</sub> gas flushing alone was effective at controlling bacterial growth for 24 h (for M2 and M3) and for 48 h (for M1) (Figure 3A, and data not shown); similarly, bacterial growth was halted for 12 h in all four milk samples (M4–M7) stored at 25°C (Figure 4A, and data not shown). Importantly, no bacterial types were clearly stimulated during the N<sub>2</sub> flushing treatments, at either considered temperature, corroborating previous observations of raw milk stored at 6 and 12°C (Munsch-Alatossava et al., 2010a; Gschwendtner et al., 2016). Compared to HT, the effectiveness of N<sub>2</sub> for the same milk sample revealed that with one exception (M1 stored at 15°C, Table 2), all ratio values determined for

N exceeded the values for HT, suggesting that altogether HT was more effective at preventing bacterial development; however, these differences were not found to be statistically significant with the REGWQ test at either considered temperature (15 or 25°C). Consequently, within a similar time frame, N<sub>2</sub> can achieve control of bacterial populations equivalent to that of the activated lactoperoxidase system (HT) (Tables 2, 3), although unlike HT, N<sub>2</sub> did not trigger any detectable bactericidal type of effect considering the original bacterial load (Figures 3B, 4B and data not shown).

As noticeable in Figures 3B, 4B, bacterial counts on MacConkey agar were lower or slightly lower at 24 h at 15°C and 12 h at 25°C than the initial counts for condition HT (this result was also observed for other samples, data not shown), illustrating the bactericidal effect of the LPs toward Gram-negative bacteria described earlier (Wolfson and Sumner, 1993; Ur-Rehman and Farkye, 2002; FAO, 2005; Seifu et al., 2005). Here, the effect of SCN<sup>-</sup> (T), sometimes clearly stimulatory, was mostly converted into high inhibition of bacterial development with the simultaneous presence of H<sub>2</sub>O<sub>2</sub> (HT) for six raw milk samples (with the exception of M1) (Tables 2, 3). Here also, the activated LPs (HT) was less effective against lactic acid bacteria, which is in agreement with previous reports (FAO, 2005).

Where refrigeration is available, the addition of H<sub>2</sub>O<sub>2</sub> is not generally recommended for raw milk; however, its use is permitted in certain applications prior to cheese making or during the preparation of modified whey, in the USA for example (Martin et al., 2014). The addition of H<sub>2</sub>O<sub>2</sub> alone (H) here did not induce high inhibition with the exception of M2 (Figure 3B). NHT yielded the lowest and the second lowest ratio in 4 and 2 cases, respectively, at 25 and 15°C (Tables 2, 3); these ratio values were very close to those obtained for HT (except for sample M3, where the difference reached 0.16). Considering the performance of HT, no synergistic effect could be induced by the additional application of N<sub>2</sub> gas flushing to the lactoperoxidase system; however, by comparing the ratios recorded for N with those of NHT, the addition of the LPs system provoked superior inhibition to that of N<sub>2</sub> flushing alone for six samples (Tables 2, 3), which altogether could suggest that the LPs-based treatment (HT) and the N<sub>2</sub> gas flushing do not target the exact same bacterial groups in raw milk.

Considering the applied treatments (with exception of T), for all bacterial populations recovered on PCA, MRS, and M17 agar, the effects were mostly bacteriostatic as a growth increase was recorded after some delay (Figures 3A, 4A, and data not shown). At both considered temperatures, a bacteriostatic type of action was also observed here with N<sub>2</sub> gas flushing (Figures 3, 4 and data not shown), which is in line with results from earlier studies when milk samples were flushed during storage between 6 and 12°C (Murray et al., 1983; Munsch-Alatossava et al., 2010a). No bactericidal type of action was noticed with N in this study, in contrast to HT. However, that N<sub>2</sub> gas flushing does not exert any bactericidal action is not that straightforward considering recent observations showing that N<sub>2</sub> gas flushing can act as a bactericidal agent for *Bacillus weihenstephanensis*, a Gram-positive representative found in pasteurized milk (Lechner et al., 1998; Munsch-Alatossava et al.,



2013), as a majority of cells were considerably damaged based on electron microscopic observations following the flushing treatment (Munsch-Alatossava and Alatossava, 2014). Most probably, our plating conditions, which did not specifically target *Bacillus* types of Gram-positive bacteria, together with the fact that *Bacillus* is usually present in low numbers in raw milk, prevented the detection of any bactericidal type of action in this study.

HT exerted rather rapid action as after 7 h at 15°C, the bacterial counts were lower than the initial counts on MacConkey agar (Figure 3B). The factor “time” seems to play also a crucial role when considering the mode of action of N<sub>2</sub> gas: for raw milk samples stored at 6°C, it appeared that phospholipase producers were excluded in a sample-dependent manner under the N<sub>2</sub> flushing treatment after 3, 7, or 11 d (Munsch-Alatossava et al., 2010b). Additionally, many days were necessary to record a bactericidal type of action on *Bacillus weihenstephanensis* under the same continuous N<sub>2</sub> gas flushing (Munsch-Alatossava and Alatossava, 2014), which altogether suggests that unlike HT (which exerts rather rapid action), the action of N<sub>2</sub> gas flushing seems to be inscribed in time.

## CONCLUSION

N<sub>2</sub>, as a non-finite resource and categorized as GRAS (Generally Regarded As Safe), also allowed in organic production systems, presents several advantages: mainly, it enables the prolonged cold

storage of milk by inhibiting spoilage bacteria without favoring well-known human pathogens or other harmful bacteria. This study shows that in the absence of cold chain facilities, N<sub>2</sub> flushing could be used within a controlled time frame (12 and 24 h at 25 and 15°C, respectively) to control bacterial growth in raw milk. Very likely, the treatment could also compensate for cold chain failures. Further studies are necessary to optimize the effectiveness of the N<sub>2</sub> gas flushing treatment combining minimal costs with maximal efficiency to offer an alternative to the use of chemicals, especially replacing the use of illegal adulterants with negative impacts on human health and the environment.

## AUTHOR CONTRIBUTIONS

PMA, IDM, and TA conceived and designed the experiments; PMA, RQ performed the experiments; JPG analyzed the data; TA, JPG contributed reagents/materials/analysis tools; PMA, TA, and JPG wrote the paper.

## ACKNOWLEDGMENTS

We gratefully acknowledge Antti Alavuotunki/Helsingin Meijeriliike Ltd for the gift of the raw milk samples. RQ was supported by the Erasmus program during March and April 2015. We thank both reviewers for their valuable comments and suggestions to improve our manuscript.

## REFERENCES

- Anonymous (2004). Regulation (EC) No 853/2004 of the European parliament and of the council of 29 April 2004 laying down specific hygiene rules for on the hygiene of foodstuffs. *Official J. Eur. Union L* 139/55. 151.
- Bafort, F., Parisi, O., Perraudin, J. P., and Jijakli, M. H. (2014). Mode of action of lactoperoxidase as related to its antimicrobial activity: a review. *Enzyme Res.* 2014:517164. doi: 10.1155/2014/517164
- Botelho, B., Reis, N., Oliveira, L. S., and Sena, M. M. (2015). Development and analytical validation of a screening method for simultaneous detection of five adulterants in raw milk using mid-infrared spectroscopy and PLS-DA. *Food Chem.* 181, 31–37. doi: 10.1016/j.foodchem.2015.02.077
- Chambers, J. V. (2002). “The microbiology of raw milk,” in *Dairy Microbiology Handbook*, ed R. K. Robinson (New York, NY: Wiley-Interscience), 39–89.
- Dechemi, S., Benjelloun, H., and Lebeault, J. M. (2005). Effect of modified atmospheres on the growth and extracellular enzymes of psychrotrophs in raw milk. *Eng. Life Sci.* 5, 350–356. doi: 10.1002/elsc.200520082
- FAO (1991). *Guidelines for the Preservation of Raw Milk by the Use of the Lactoperoxidase System*. CAC/GL13–1991.
- FAO (2005). “Benefits and potential risks of the lactoperoxidase system of raw milk preservation,” in *Report of an FAO/WHO Technical Meeting* (Rome: FAO Headquarters).
- FAO (2012). Available online at: [www.fao.org/save-food/resources/keyfindings/infographics/dairy/en](http://www.fao.org/save-food/resources/keyfindings/infographics/dairy/en)
- Frank, J. F., and Hassan, A. N. (2002). “Microorganisms associated with milk,” in *Encyclopedia of Dairy Sciences*, Vol. 3, eds H. Roginski, J.W. Fuquay and P.F. Fox (London: Academic Press), 1786–1796.
- Fweja, L. W. T., Lewis, M. J., and Grandison, A. S. (2008). Challenge testing the lactoperoxidase system against a range of bacteria using different activation agents. *J. Dairy Sci.* 91, 2566–2574. doi: 10.3168/jds.2007-0322
- Gschwendtner, S., Alatossava, T., Kublik, S., Fuka, M. M., Schloter, M., and Munsch-Alatossava, P. (2016). N<sub>2</sub> gas flushing alleviates the loss of bacterial diversity and inhibits psychrotrophic *Pseudomonas* during the cold storage of bovine raw milk. *PLoS ONE* 11:e014605. doi: 10.1371/journal.pone.0146015
- Gustavsson, J., Cederberg, C., Sonesson, U., Van Otterdijk, R., and Meybeck, A. (2011). “Global food losses and food waste,” in *Save Food at Interpack* (Düsseldorf; Rome: Food and Agricultural Organisation of the United Nations), 38.
- Hsu, J. C. (1996). *Multiple Comparisons: Theory and Methods*. London: Chapman and Hall.
- Korhonen, H., and Marnila, P. (2002). “Lactoferrin,” in *Encyclopedia of Dairy Sciences*, Vol. 3, eds H. Roginski, J. W. Fuquay and P. F. Fox (London: Academic Press), 1946–1950.
- Lechner, S., Mayr, R., Francis, K. P., Prüß, B. M., Kaplan, T., Wießner-Gunkel, et al. (1998). *Bacillus weihenstephanensis* sp. nov. is a new psychrotolerant species of the *Bacillus cereus* group. *Int. J. Syst. Bacteriol.* 48, 1373–1382.
- Marnila, P., and Korhonen, H. (2002). “Immunoglobulins,” in *Encyclopedia of Dairy Sciences*, Vol. 3, eds H. Roginski, J. W. Fuquay and P. F. Fox (London: Academic Press), 1050–1056.
- Martin, N. H., Friedlander, A., Mok, A., Kent, D., Wiedmann, M., and Boor, K. J. (2014). Peroxide test strips detect added hydrogen peroxide in raw milk at levels affecting bacterial load. *J. Food Prot.* 77, 1809–1813. doi: 10.4315/0362-028X.JFP-14-074
- Munsch-Alatossava, P., and Alatossava, T. (2007). Antibiotic resistance of raw-milk associated psychrotrophic bacteria. *Microbiol. Res.* 162, 115–123. doi:10.1016/j.micres.2006.01.015
- Munsch-Alatossava, P., and Alatossava, T. (2014). Nitrogen gas flushing can be bactericidal: the temperature-dependent destiny of *Bacillus weihenstephanensis* KBAB4 under a pure N<sub>2</sub> atmosphere. *Front. Microbiol.* 5:619. doi: 10.3389/fmicb.2014.00619
- Munsch-Alatossava, P., Gauchi, J. P., Chamlagain, B., and Alatossava, T. (2012a). Trends of antibiotic resistance in mesophilic and psychrotrophic bacterial

- populations during cold storage of raw milk. *ISRN Microbiol.* 2012:918208. doi:105402/2012/918208
- Munsch-Alatossava, P., Ghafar, A., and Alatossava, T. (2013). Potential of nitrogen gas (N<sub>2</sub>) flushing to extend the shelf life of cold stored pasteurised milk. *Int. J. Mol. Sci.* 14, 5668–5685. doi: 10339/ijms14035668
- Munsch-Alatossava, P., GURSOY, O., and Alatossava, T. (2010a). Potential of nitrogen gas (N<sub>2</sub>) to control psychrotrophs and mesophiles in raw milk. *Microbiol. Res.* 165, 122–132. doi: 10.1016/j.micres.2009.02.002
- Munsch-Alatossava, P., GURSOY, O., and Alatossava, T. (2010b). Exclusion of phospholipases (PLs)-producing bacteria in raw milk flushed with nitrogen gas (N<sub>2</sub>). *Microbiol. Res.* 165, 61–65. doi: 10.1016/j.micres.2008.07.001
- Munsch-Alatossava, P., GURSOY, O., and Alatossava, T. (2011). Improved storage of cold raw milk by continuous flushing of N<sub>2</sub> gas separated from compressed air: a pilot scale study. *J. Food Process Technol.* 1, 1–4. doi: 10.4172/2157-7110.1000101
- Munsch-Alatossava, P., Ikonen, V., Alatossava, T., and Gauchi, J. P. (2012b). “Trends of antibiotic resistance (AR) in mesophilic and psychrotrophic bacterial populations during cold storage of raw milk, produced by organic and conventional farming systems,” in *Antibiotic Resistant Bacteria, A Continuous Challenge in the New Millenium*, ed M. Pana (Rijeka: Intech), 105–124.
- Murray, S. K., Kwan, K. K. H., Skura, B. J., and McKellar, R. C. (1983). Effect of nitrogen flushing on the production of proteinase by psychrotrophic bacteria in raw milk. *J. Food Sci.* 48, 1166–1169
- Reiter, B., and Härnuly, G. (1984). Lactoperoxidase antibacterial system: natural occurrence, biological functions and practical applications. *J. Food Prot.* 47, 724–732.
- Seifu, E., Buys, E. M., and Donkin, E. F. (2005). Significance of the lactoperoxidase system in the dairy industry and its potential applications: a review. *Trends Food Sci. Technol.* 16, 137–154. doi: 10.1016/j.tifs.2004.11.002
- Ur-Rehman, S., and Farkye, N. Y. (2002). “Lactoperoxidase,” in *Encyclopedia of Dairy Sciences, Vol. 2*, eds H. Roginski, J. W. Fuquay and P. F. Fox (London: Academic Press), 938–941.
- Wolfson, L., and Sumner, S. S. (1993). Antibacterial activity of the lactoperoxidase system: a review. *J. Food Prot.* 56, 887–892.

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Munsch-Alatossava, Quintyn, De Man, Alatossava and Gauchi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.