



Endocannabinoid System and Migraine Pain: An Update

Rosaria Greco^{1*}, Chiara Demartini¹, Anna M. Zanaboni^{1,2}, Daniele Piomelli³ and Cristina Tassorelli^{1,2}

¹ Laboratory of Neurophysiology of Integrative Autonomic Systems, Headache Science Centre, IRCCS Mondino Foundation, Pavia, Italy, ² Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy, ³ Department of Anatomy and Neurobiology, University of California, Irvine, Irvine, CA, United States

OPEN ACCESS

Edited by:

Fabrizio A. Pamplona,
Entourage Phytolab, Brazil

Reviewed by:

Gabriela Rodriguez-Manzo,
Centro de Investigación y de Estudios
Avanzados del Instituto Politécnico
Nacional (CINVESTAV-IPN), Mexico
Mariana Spetea,
University of Innsbruck, Austria

*Correspondence:

Rosaria Greco
rosaria.greco@mondino.it

Specialty section:

This article was submitted to
Neuropharmacology,
a section of the journal
Frontiers in Neuroscience

Received: 21 December 2017

Accepted: 05 March 2018

Published: 19 March 2018

Citation:

Greco R, Demartini C, Zanaboni AM,
Piomelli D and Tassorelli C (2018)
Endocannabinoid System and
Migraine Pain: An Update.
Front. Neurosci. 12:172.
doi: 10.3389/fnins.2018.00172

The trigeminovascular system (TS) activation and the vasoactive release from trigeminal endings, in proximity of the meningeal vessels, are considered two of the main effector mechanisms of migraine attacks. Several other structures and mediators are involved, however, both upstream and alongside the TS. Among these, the endocannabinoid system (ES) has recently attracted considerable attention. Experimental and clinical data suggest indeed a link between dysregulation of this signaling complex and migraine headache. Clinical observations, in particular, show that the levels of anandamide (AEA)—one of the two primary endocannabinoid lipids—are reduced in cerebrospinal fluid and plasma of patients with chronic migraine (CM), and that this reduction is associated with pain facilitation in the spinal cord. AEA is produced on demand during inflammatory conditions and exerts most of its effects by acting on cannabinoid (CB) receptors. AEA is rapidly degraded by fatty acid amide hydrolase (FAAH) enzyme and its levels can be modulated in the peripheral and central nervous system (CNS) by FAAH inhibitors. Inhibition of AEA degradation via FAAH is a promising therapeutic target for migraine pain, since it is presumably associated to an increased availability of the endocannabinoid, specifically at the site where its formation is stimulated (e.g., trigeminal ganglion and/or meninges), thus prolonging its action.

Keywords: migraine, inflammation, trigeminal hyperalgesia, endocannabinoid system, FAAH inhibitors

INTRODUCTION

Migraine is one of the most disabling painful conditions and a very common disorder (Global Burden of Disease, 2015). Although the pathophysiology of migraine is still largely elusive, the trigeminovascular system (TS) activation and the neurogenic inflammation of the dura mater are widely recognized as two key mechanisms underlying migraine attacks (Moskowitz, 1993). TS activation causes neuropeptide release from trigeminal endings in proximity of the meningeal vessels. Meningeal release of mediators produces peripheral sensitization, which is aggravated by central sensitization when the attacks recur more frequently. Calcitonin gene-related peptide (CGRP) and other inflammatory mediators, released from sensory nerve terminals (Moskowitz, 1993), irritate and further dilate blood vessels, thus resulting in the release of additional neuropeptides from the sensory neurons and an increase of pain impulses that are transmitted to the nucleus trigeminalis caudalis (NTC). The activated NTC relays in turn pain signals to higher brain centers, including thalamus and cortex. In this circuitry, another interesting player is nitric oxide (NO), which contributes to the perivascular sensory afferent nerve fibers

activation in the meninges and to neuropeptides release or NO formation by neuronal NO synthase (nNOS) (Messlinger et al., 2000; Alaşehirli et al., 2013; Ramachandran et al., 2014). Evidence suggests that the origin of migraine attacks is the interaction of internal or external triggers with dysfunctional central structures (brainstem, thalamus) involved in the transmission and regulation of pain sensation (Goadsby, 2002; Knight et al., 2005; Coppola et al., 2013).

Current standards of care for migraine have a moderate effectiveness at best and, in some cases, limited tolerability. Specifically, prophylactic treatments (beta blockers, antiepileptic drugs) may induce weight gain, depression, behavioral or cognitive disturbances. Triptans, 5-HT₁-Receptor agonists, used for acute treatment, may cause a serious long-term side effects such as chest pain, neck and limbs with paresthesias and hot or cold sensations. Analgesics combinations and non-steroidal anti-inflammatory drugs, for acute migraine can lead to gastrointestinal and cardio-renal side effects (Antonaci et al., 2016). The endocannabinoid system (ES) has recently received attention in regard to pain control, after the availability of probes capable of modulating its activity via the interaction with endocannabinoid catabolic enzymes (Chiou et al., 2013). In this review, we summarize results collected in studies aimed at evaluating the role of the endocannabinoids (ECs) in migraine, with a specific focus on fatty acid amide hydrolase (FAAH) inhibitors.

ENDOCANNABINOID SIGNALING

EC signaling in the nervous system is different from those of the classic neurotransmission systems, where the depolarization of the presynaptic neuron causes neurotransmitters release which in turn activates their receptors on the postsynaptic neuron. The ECs act as retrograde or local neurotransmitters, and are produced and released from neurons upon demand. They bind to CB₁-type cannabinoid (CB₁) receptors, which are localized on presynaptic terminals of excitatory and inhibitory neurons throughout the brain and spinal cord (Alger and Kim, 2011). CB₁ receptors are seven trans-membrane domain proteins that belong to the rhodopsin family of G protein-coupled receptors, specifically those of the Gi/o family (GPCRs). Recent crystallographic studies reported that extracellular surface of CB₁ receptor is different from other lipid-activated GPCRs with a critical part of the ligand binding pocket (Hua et al., 2016; Shao et al., 2016). CB₁ receptors are found in neuroanatomical regions involved in pain processing, and inhibit the release of neurotransmitters such as γ -aminobutyric acid, glutamate, dopamine, noradrenaline and acetylcholine (Katona and Freund, 2008). Though abundant in neurons of the central nervous system (CNS), CB₁ receptors are also expressed in peripheral neurons and many non-neural cells (Kendall and Yudowski, 2017). A second cannabinoid receptor subtype, CB₂, is found primarily in immune cells (Gerdeman et al., 2002). Furthermore AEA and 2-arachidonoylglycerol (2-AG), the best characterized ECs, are produced in structures involved in nociception, such as the skin, dorsal root ganglia,

spinal cord, periaqueductal gray matter (PAG), and rostral ventromedial medulla (RVM) (Katona and Freund, 2008). Through activation of CB₁ receptors, AEA and 2-AG can influence a variety of physiological processes, including energy balance, cognition and pain (Bellocchio et al., 2008; Kano et al., 2009).

In neurons, as in other cells, the ECs are not stored in vesicles but are enzymatically produced upon demand from membrane glycerophospholipid precursors. Enzymes involved in AEA and 2-AG formation are N-acylphosphatidylethanolamine-phospholipase D (NAPE-PLD) and diacylglycerol lipase (DGL), respectively (Bisogno et al., 2003; Okamoto et al., 2007). However, other pathways through which AEA can be produced have been described (Liu et al., 2006; Jin et al., 2007). Moreover, several enzymes involved in ECs biosynthesis, such as NAPE-PLD, give rise not only to AEA but also to structurally similar lipid messengers that do not bind and activate CB₁, i.e., oleoylethanolamide (OEA) and palmitoylethanolamide (PEA) (Gaetani et al., 2010). AEA acts primarily on CB₁ receptors, though pharmacological actions on other receptors, such as transient receptor potential (TRP) V₁, have been described (Puente et al., 2011), TRPV₂, TRPA₁, and TRPM₈ (Di Marzo and De Petrocellis, 2010).

2-AG production also occurs via multiple biosynthetic pathways, in which diacylglycerol (DAG), produced by the action of either phospholipase C (PLC) or phosphatidic acid phosphohydrolase, acts as a common precursor. DAG is transformed into 2-AG by DGL; alternatively, phospholipase A₁ may convert phosphatidyl-inositol into lyso-phosphatidyl-inositol, which may be transformed to 2-AG by PLC.

The ECs are quickly deactivated by uptake into cells followed by intracellular hydrolysis (Urquhart et al., 2015). Transporter proteins remove AEA from the extracellular space; successively AEA is mostly degraded by FAAH, releasing arachidonic acid (AA) and ethanolamine. 2-AG is hydrolyzed mainly by the serine hydrolase, monoacylglycerol lipase (MGL), which produces AA and glycerol. However, it may be also degraded by α,β -hydrolase-6 or converted to bioactive oxygenated products by COX₂. Thus, the enzymes responsible for the biosynthetic, as well as degradative pathways are essential in the regulation and modulation of EC levels in the CNS. Moreover, differential cellular distribution of the synthesizing and degrading enzymes may control of EC activity. Thus, selective pharmacological or genetic manipulations of FAAH and MGL activities can be used to evaluate the functions of each EC in animal model.

RELATIONSHIP BETWEEN ES DYSREGULATION AND MIGRAINE: HUMAN AND EXPERIMENTAL STUDIES

The ES may modulate the cerebrovascular tone, through interaction with serotonergic system, NO synthesis, and neuropeptides release (Pertwee, 2001), neurotransmitters that play a crucial role in migraine pathogenesis. CB₁ receptors have been localized in potential generators of migraine pain, like PAG, RVM, and NTC (Moldrich and Wenger, 2000). There are reports

that frequency of migraine headache may decrease in persons using medical cannabis (Rhyne et al., 2016). ECs may interact with and modulate several pathways related to migraine, such as opioids, or involved in the mechanism of action of anti-migraine drugs such as triptans (Akerman et al., 2013; Baron, 2015). AEA and other CB agonists have also been demonstrated to inhibit effects on serotonin type 3 receptors, which provide yet another effect when considering that nausea and vomiting are frequent and bothersome accompaniments of migraine (Fan, 1995; Park et al., 2008). CB agonists inhibit the serotonin-induced current in a concentration dependent manner in the rat nodose ganglion neurons by 5-HT₃ receptor ion-channel (Fan, 1995). Moreover, they may also act on brain areas involved in emesis, such as the dorsal motor nucleus of the vagus (Van Sickle et al., 2001), where there is a high density of 5-HT₃ receptors (Miquel et al., 2002). 5-HT₃ inhibition can modulate neurotransmitters, including dopamine, GABA, substance P, and acetylcholine. The anti-migraine effects of the ES are not fully known, although some hypotheses were proposed. **Table 1** shows the potential modulatory effects of ECs on migraine pain.

Clinical observations show that women migraine without aura or episodic tension-type headache have increased FAAH and endocannabinoid membrane transporter (EMT) activities in platelets, which is consistent with reduced AEA levels (Cupini et al., 2006). In addition, women with episodic migraine have increased CB₁ receptor binding during the interictal period, as assessed by positron emission tomography; this increase is especially evident in brain regions that exert top-down influences to modulate pain (Van der Schueren et al., 2012). Variants in the CB₁ receptor gene increase the risk of migraine attack with nausea in life stress exposed subjects (Juhász et al., 2017). Recently Gouveia-Figueira et al. (2017) failed to detect significant changes in the plasma levels of AEA and other fatty acid ethanolamides between patients with episodic migraine and controls. These contrasting findings may be related to higher inter-subject variability of EC levels in the evaluated cohorts or to a different migraine load on the populations investigated.

More consistent are the findings regarding the involvement of the ES in chronic migraine (CM). Subjects with CM with and without medication overuse headache (MOH) showed reduced activities of FAAH and EMT in platelets when compared to either controls or episodic migraine (Cupini et al., 2008). In another study, 2-AG and AEA platelet levels were significantly lower in MOH and CM patients compared to controls, without significant

differences between the two patient groups (Rossi et al., 2008). These findings suggest an adaptive behavior induced by chronic headache *per se*, while medication overuse is apparently not related with EC activity. Interestingly, serotonin levels were reduced in the MOH and CM patients, with lower values detected in females as compared to males (Rossi et al., 2008) and that serotonin levels were also associated with 2-AG tone, with a higher correlation coefficient for MOH patients. This latter finding suggests a possible role for 2-AG, together with serotonin, in the “addiction” aspect of MOH. In this frame, it is worth mentioning that successful detoxification of MOH subjects is accompanied by a reduction in FAAH activity in platelets. This biochemical change is associated with the normalization of neurophysiological parameters that indicated a status of sensitization of the pain pathways (Perrotta et al., 2012). A reduction in AEA and an increase in PEA levels was also found in the cerebrospinal fluid of both CM and MOH patients (Sarchielli et al., 2007), pointing to a central alteration of ES in these subjects.

Inflammation and nerve injury cause changes in local AEA levels (Jhaveri et al., 2007). As mentioned before, AEA is produced on demand during inflammatory conditions and it is rapidly degraded by FAAH activity. Thus, AEA tone can be modulated by FAAH activity in both periphery and CNS. Increased activation of the TS may theoretically lead to reduced levels of AEA, which may, in turn, lead to an increased CGRP and NO release. AEA indeed inhibits the neurogenic dural vasodilatation, as well as CGRP-induced and NO-induced dural vessel dilation (Akerman et al., 2004). The CB₁ receptor antagonist, AM251, reversed this inhibitory activity, suggesting that CB₁ receptors may be implicated in the relationship between headache and dural blood vessel dilation and migraine mediators. Cortical spreading depression (CSD) is a self-propagating wave of neuronal hyperexcitability that has a role in migraine (Goadsby, 2007). WIN55,212-2, a CB₁ receptor agonist, inhibited the amplitude, duration and velocity of CSD propagation, while JWH 133, a CB₂ receptor agonist, was devoid of any effect (Kazemi et al., 2012).

The trigeminal firing in the trigeminocervical complex induced by AEA inhibition is reversed after CB₁ receptor antagonism, thus suggesting that the central effects of AEA are principally CB₁-mediated (Akerman et al., 2007). CB₁ receptor activation in the ventrolateral PAG, obtained with the administration of WIN55,212-2, attenuates the activity evoked by dural stimulation in A-fiber neurons and the basal spontaneous

TABLE 1 | Potential effects of endocannabinoids on migraine pain.

Target	Effects	References
Trigeminovascular activation	Substance P ↓ CGRP/nitric oxide ↓ Cyclooxygenase ↓ PGE-2 synthesis ↓ glutamate release ↓	Pertwee, 2001; Akerman et al., 2004; Sarchielli et al., 2007; La Rana et al., 2008; Chiou et al., 2013
Serotonergic system	Serotonin release ↓ platelets aggregation ↓ 5-HT _{2A} ↑	Volfe et al., 1985; Ohuoha et al., 1994; Boger et al., 1998; Rossi et al., 2008; Parker et al., 2011; Mendiguren et al., 2018
Brainstem	NF-κB activation ↓ kynurenine pathway modulation	Kelly and Chapman, 2001; Nagy-Grócz et al., 2016
Hypothalamus	Glutamate release ↓	Di et al., 2005
Periaqueductal gray	Proenkephalin expression ↑	Manzanares et al., 1998

activity in the trigeminocervical complex of rodent. These findings suggest that, in the brainstem, ECs may provide to descending modulation upon basal trigeminovascular neuronal tone and A-fiber dural-nociceptive responses, (Akerman et al., 2013). Changes in FAAH and MGL activities were found in the brainstem and hypothalamus of rats treated with nitroglycerin (NTG) (Greco et al., 2010b), a recognized animal model of migraine (Buzzi and Tassorelli, 2010). NTG in rat causes an increased sensitivity to nociceptive tests and c-fos protein expression in brain areas nuclei involved in migraine pain transmission, such as NTC (Greco et al., 2011a). The use of this model by us and other groups has allowed the in-depth exploration of the mechanisms underlying the modulation of the ECs and the nociceptive activation of the TS during a migraine attack. In particular, we reported an increased FAAH activity in the hypothalamus and in the medulla area, where NTC neurons are located, and an up-regulation of CB₁ receptor binding sites in the same areas (Greco et al., 2010b), suggesting a key role of AEA in the cephalic pain.

Our findings also show that AEA pretreatment significantly reduces NTG-induced behavioral nocifensive and NTG-induced neuronal activation in the NTC (Greco et al., 2011a); moreover, AEA may modulate central sensitization through TRPV1, COX2 expression and NF- κ B inhibition in NTC (Nagy-Grócz et al., 2016). The CB₂ receptors activation in pain modulation has been considered in the past, showing analgesic activity in several models of pain (Nackley et al., 2003, 2004; Quartilho et al., 2003). In our migraine model, we have also shown that CB₂ receptor activation significantly decreases nocifensive behavior of rats made hyperalgesic by NTG (Greco et al., 2014). Likewise, MGL inhibition, and the subsequent increase in central and/or peripheral levels of 2-AG, reduces NTG-induced hyperalgesia at the nociceptive tests, and attenuates c-Fos protein expression in brain areas implicated in the transmission or integration of cephalic pain (Greco et al., 2017).

RECENT ADVANCES ON FAAH INHIBITION IN MIGRAINE PAIN

Though the analgesic effects of cannabinoids are fairly well established, their use in therapy remains limited by their psychoactive properties (Borgelt et al., 2013). Recent safety concerns about FAAH inhibitors turned out to be ungrounded, and due to off-target effects. Clearly, the successful development of compounds that modulate ECs tone for the pain relief in humans will hinge on the ability to separate psychotropic effects from therapeutic ones, and to control for potential off-target interactions. Positive allosteric modulation of CB₁ receptor signaling may represent a safe analgesic alternative strategy that lacks tolerance, dependence and abuse liability (Khurana et al., 2017; Slivicki et al., 2017). Several studies show that also increasing ECs levels through the inhibition of catabolic enzymes, FAAH in particular, would decrease cannabimimetic side effects (Piomelli et al., 2006; Booker et al., 2012).

Besides AEA, FAAH degrades other fatty acid amides, which have several biological functions and mechanisms of action (Ahn

et al., 2008). FAAH is contained in intracellular membranes of postsynaptic somata and dendrites of the mammalian brain (Gulyas et al., 2004). In many cerebral structures FAAH and CB₁ receptors cellular co-localization in cell bodies or dendrites in proximity of CB₁-expressing fibers (Egertová et al., 1998). Manipulations of full-length and transmembrane-truncated FAAH variants have offered a characterization of mechanisms of action (McKinney and Cravatt, 2005). In particular, these studies showed that, unlike most serine hydrolases, which use a histidine residue as a catalytic base, FAAH recruits a lysine to hydrolyze both amides and esters at equivalent rates (Patricelli and Cravatt, 1999). Numerous FAAH inhibitors have been developed and tested in animal models of disease (Jayamanne et al., 2006; Kinsey et al., 2009). In particular, the FAAH inhibition induces anti-inflammatory effects *in vivo* (Jayamanne et al., 2006; Booker et al., 2012; Wilkerson et al., 2017). In addition, mutant mice for FAAH enzyme in non-neuronal cells, but with FAAH activity conserved in peripheral and central neurons, have a phenotype in which basal nociceptive transmission is connected to the reduced responsiveness to pro-inflammatory mediators (Cravatt et al., 2004). Researchers suggest that AEA regulates nociceptive transmission primarily at the peripheral level (Calignano et al., 1998; Clapper et al., 2010; Piomelli and Sasso, 2014).

Numerous studies have shown that FAAH inhibition causes analgesia and reduces inflammation in animal models of acute inflammatory pain (Kinsey et al., 2010; Lodola et al., 2015; Nasirinezhad et al., 2015), but there is little information on their effects in migraine. Recently, it was reported that AEA modulates the analgesic activity in the orofacial area and that endomorphin-2-induced antinociception is mediated by μ and CB₁ receptors (Zubrzycki et al., 2017). Nozaki et al. (2015) demonstrated that NTG-induced mechanical allodynia and c-Fos protein in the NTC is abolished in FAAH-deficient mice or after URB597 treatment, a global FAAH inhibitor, via maintenance of central and peripheral AEA levels. When considering that NTG is thought to activate meningeal trigeminovascular terminals via the local NO formation (Reuter et al., 2001; Greco et al., 2011b), it is probable that URB597 interferes with this mechanism of peripheral sensitization. Accordingly, we have shown that a peripherally restricted FAAH inhibitor, the compound URB937, inhibits NTG-induced nocifensive behaviors (plantar and orofacial formalin test, tail flick test), neuronal activation in the NTC and locus coeruleus (Greco et al., 2015). In agreement with these data, URB937 decreases the c-Fos expression induced by plantar formalin injection in spinal cord regions involved in nociceptive processing by the CB₁ receptors (Clapper et al., 2010).

Thus, since URB937 acts only peripherally, it seems reasonable to hypothesize that its mechanism of action relies on the maintenance of higher levels of AEA released by nervous terminal located in the injured peripheral tissues (hindpaw, upper lip, tail) (Agarwal et al., 2007) or in the dura, with consequent CB₁ receptor activation in trigeminovascular endings. An additional mechanism, is probably represented by the blockade of NTG-induced inflammatory pathway mediated by NO in dura mater and/or trigeminal ganglia. In agreement

with this hypothesis, *in vitro* studies have shown that increased AEA tone, through the inhibition of its degradation or uptake, decreases the cytokines and NO levels (Correa et al., 2009, 2010).

OUTLOOK

Pain is a heterogeneous condition and it should be treated as such. With its lack of sensitivity to standard analgesic medications (Ong and De Felice, 2017), migraine pain is a case in point and—perhaps better than most other forms of pain—underscores the need for tailored therapies. The human data and preclinical studies reviewed here confirm the importance of FAAH-regulated AEA signaling in the processing of nociceptive signals outside the CNS (Greco et al., 2010a; Piomelli and Sasso, 2014) and specifically point to peripheral FAAH inhibition as a possible therapeutic opportunity for migraine pain. Future

experiments should be aimed at unlocking the precise cellular mechanisms and neural circuits through which peripheral FAAH blockade exerts its analgesic effects in migraine pain, further exploring the ground for potential clinical trials.

AUTHOR CONTRIBUTIONS

RG: designed this review; CD and AZ: contributed to cited data from our group; DP and CT: revised the final version of the manuscript.

FUNDING

Some of our data on FAAH inhibition reported were supported by a grant from the Italian Ministry of Health to C. Mondino National Neurological Institute (RF2013-02355704).

REFERENCES

- Agarwal, N., Pacher, P., Tegeder, I., Amaya, F., Constantin, C. E., Brenner, G. J., et al. (2007). Cannabinoids mediate analgesia largely via peripheral type 1 cannabinoid receptors in nociceptors. *Nat. Neurosci.* 10, 870–879. doi: 10.1038/nn1916
- Ahn, K., McKinney, M. K., and Cravatt, B. F. (2008). Review enzymatic pathways that regulate endocannabinoid signaling in the nervous system. *Chem. Rev.* 108, 1687–1707. doi: 10.1021/cr0782067
- Akerman, S., Holland, P. R., and Goadsby, P. J. (2007). Cannabinoid (CB1) receptor activation inhibits trigeminovascular neurons. *J. Pharmacol. Exp. Ther.* 320, 64–71. doi: 10.1124/jpet.106.106971
- Akerman, S., Holland, P. R., Lasalandra, M. P., and Goadsby, P. J. (2013). Endocannabinoids in the brainstem modulate dural trigeminovascular nociceptive traffic via CB1 and “triptan” receptors: implications in migraine. *J. Neurosci.* 33, 14869–14877. doi: 10.1523/JNEUROSCI.0943-13.2013
- Akerman, S., Kaube, H., and Goadsby, P. J. (2004). Anandamide is able to inhibit trigeminal neurons using an *in vivo* model of trigeminovascular-mediated nociception. *J. Pharmacol. Exp. Ther.* 309, 56–63. doi: 10.1124/jpet.103.059808
- Alaşehirli, B., Akçali, A., Demiryürek, A. T., Özel, A., Erdal, M. E., and Neyal, M. (2013). Lack of association between the C276T polymorphism of the neuronal nitric oxide synthase gene and migraine. *Int. J. Neurosci.* 123, 50–54. doi: 10.3109/00207454.2012.724743
- Alger, B. E., and Kim, J. (2011). Supply and demand for endocannabinoids. *Trends Neurosci.* 34, 304–315. doi: 10.1016/j.tins.2011.03.003
- Antonaci, F., Ghiotto, N., Wu, S., Pucci, E., and Costa, A. (2016). Recent advances in migraine therapy. *Springerplus* 5:637. doi: 10.1186/s40064-016-2211-8
- Baron, E. P. (2015). Comprehensive review of medicinal Marijuana, Cannabinoids, and therapeutic implications in medicine and headache: what a long strange trip it's been. *Headache* 55, 885–916. doi: 10.1111/head.12570
- Bellocchio, L., Cervino, C., Pasquali, R., and Pagotto, U. (2008). The endocannabinoid system and energy metabolism. *J. Neuroendocrinol.* 20, 850–807. doi: 10.1111/j.1365-2826.2008.01728.x
- Bisogno, T., Howell, F., Williams, G., Minassi, A., Cascio, M. G., Ligresti, A., et al. (2003). Cloning of the first sn1-DAG lipases points to the spatial and temporal regulation of endocannabinoid signaling in the brain. *J. Cell Biol.* 163, 463–468. doi: 10.1083/jcb.200305129
- Boger, D. L., Patterson, J. E., and Jin, Q. (1998). Structural requirements for 5-HT_{2A} and 5-HT_{1A} serotonin receptor potentiation by the biologically active lipid oleamide. *Proc. Natl. Acad. Sci. U.S.A.* 95, 4102–4107. doi: 10.1073/pnas.95.8.4102
- Booker, L., Kinsey, S. G., Abdullah, R. A., Blankman, J. L., Long, J. Z., Ezzili, C., et al. (2012). The fatty acid amide hydrolase (FAAH) inhibitor PF-3845 acts in the nervous system to reverse LPS-induced tactile allodynia in mice. *Br. J. Pharmacol.* 165, 2485–2496. doi: 10.1111/j.1476-5381.2011.01445.x
- Borgelt, L. M., Franson, K. L., Nussbaum, A. M., and Wang, G. S. (2013). The pharmacologic and clinical effects of medical cannabis. *Pharmacotherapy* 33, 195–209. doi: 10.1002/phar.1187
- Buzzi, M. G., and Tassorelli, C. (2010). Experimental models of migraine. *Handb. Clin. Neurol.* 97, 109–123. doi: 10.1016/S0072-9752(10)97008-5
- Calignano, A., La Rana, G., Giuffrida, A., and Piomelli, D. (1998). Control of pain initiation by endogenous cannabinoids. *Nature* 394, 277–281. doi: 10.1038/28393
- Chiou, L. C., Hu, S. S., and Ho, Y. C. (2013). Targeting the cannabinoid system for pain relief? *Acta Anaesthesiol. Taiwan* 51, 161–170. doi: 10.1016/j.aat.2013.10.004
- Clapper, J. R., Moreno-Sanz, G., Russo, R., Guijarro, A., Vacondio, F., Duranti, A., et al. (2010). Anandamide suppresses pain initiation through a peripheral endocannabinoid mechanism. *Nat. Neurosci.* 13, 1265–1270. doi: 10.1038/nn.2632
- Coppola, G., Di Lorenzo, C., Schoenen, J., and Pierelli, F. (2013). Habituation and sensitization in primary headaches. *J. Headache Pain* 14:65. doi: 10.1186/1129-2377-14-65
- Correa, F., Docagne, F., Mestre, L., Clemente, D., Hernangómez, M., Loria, F., et al. (2009). A role for CB2 receptors in anandamide signalling pathways involved in the regulation of IL-12 and IL-23 in microglial cells. *Biochem. Pharmacol.* 77, 86–100. doi: 10.1016/j.bcp.2008.09.014
- Correa, F., Hernangómez, M., Mestre, L., Loria, F., Spagnolo, A., Docagne, F., et al. (2010). Anandamide enhances IL-10 production in activated microglia by targeting CB(2) receptors: roles of ERK1/2, JNK, and NF-kappaB. *Glia* 58, 135–147. doi: 10.1002/glia.20907
- Cravatt, B. F., Saghatelian, A., Hawkins, E. G., Clement, A. B., Bracey, M. H., and Lichtman, A. H. (2004). Functional disassociation of the central and peripheral fatty acid amide signaling systems. *Proc. Natl. Acad. Sci. U.S.A.* 101, 10821–10826. doi: 10.1073/pnas.0401292101
- Cupini, L. M., Bari, M., Battista, N., Argirò, G., Finazzi-Agrò, A., Calabresi, P., et al. (2006). Biochemical changes in endocannabinoid system are expressed in platelets of female but not male migraineurs. *Cephalalgia* 26, 277–281. doi: 10.1111/j.1468-2982.2005.01031.x
- Cupini, L. M., Costa, C., Sarchielli, P., Bari, M., Battista, N., Eusebi, P., et al. (2008). Degradation of endocannabinoids in chronic migraine and medication overuse headache. *Neurobiol. Dis.* 30, 186–189. doi: 10.1016/j.nbd.2008.01.003
- Di, S., Boudaba, C., Popescu, I. R., Weng, F. J., Harris, C., Marcheselli, V. L., et al. (2005). Activity-dependent release and actions of endocannabinoids in the rat hypothalamic supraoptic nucleus. *J. Physiol.* 569, 751–760. doi: 10.1113/jphysiol.2005.097477
- Di Marzo, V., and De Petrocellis, L. (2010). Endocannabinoids as regulators of transient receptor potential (TRP) channels: a further opportunity to develop new endocannabinoid-based therapeutic drugs. *Curr. Med. Chem.* 17, 1430–1449. doi: 10.2174/092986710790980078

- Egertová, M., Giang, D. K., Cravatt, B. F., and Elphick, M. R. (1998). A new perspective on cannabinoid signalling: complementary localization of fatty acid amide hydrolase and the CB1 receptor in rat brain. *Proc. Biol. Sci.* 265, 2081–2085. doi: 10.1098/rspb.1998.0543
- Fan, P. (1995). Cannabinoid agonists inhibit the activation of 5-HT3 receptors in rat nodose ganglion neurons. *J. Neurophysiol.* 73, 907–910. doi: 10.1152/jn.1995.73.2.907
- Gaetani, S., Fu, J., Cassano, T., Dipasquale, P., Romano, A., Righetti, L., et al. (2010). The fat-induced satiety factor oleoylethanolamide suppresses feeding through central release of oxytocin. *J. Neurosci.* 30, 8096–8101. doi: 10.1523/JNEUROSCI.0036-10.2010
- Gerdeman, G. L., Ronesi, J., and Lovinger, D. M. (2002). Postsynaptic endocannabinoid release is critical to long-term depression in the striatum. *Nat. Neurosci.* 5, 446–451. doi: 10.1038/nn832
- Global Burden of Disease (2015). Disease and Injury Incidence and Prevalence Collaborators 2016. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 388, 1545–1602. doi: 10.1016/S0140-6736(16)31678-6
- Goadsby, P. J. (2002). Neurovascular headache and a midbrain vascular malformation: evidence for a role of the brainstem in chronic migraine. *Cephalalgia* 22, 107–111. doi: 10.1046/j.1468-2982.2002.00323.x
- Goadsby, P. J. (2007). Recent advances in understanding migraine mechanisms, molecules and therapeutics. *Trends Mol. Med.* 13, 39–44. doi: 10.1016/j.molmed.2006.11.005
- Gouveia-Figueira, S., Goldin, K., Hashemian, S. A., Lindberg, A., Persson, M., Nording, M. L., et al. (2017). Plasma levels of the endocannabinoid anandamide, related N-acyl ethanolamines and linoleic acid-derived oxylipins in patients with migraine. *Prostaglandins Leukot. Essent. Fatty Acids* 120, 15–24. doi: 10.1016/j.plefa.2017.04.005
- Greco, R., Bandiera, T., Mangione, A. S., Demartini, C., Siani, F., Nappi, G., et al. (2015). Effects of peripheral FAAH blockade on NTG-induced hyperalgesia—evaluation of URB937 in an animal model of migraine. *Cephalalgia* 35, 1065–1076. doi: 10.1177/0333102414566862
- Greco, R., Demartini, C., Zanaboni, A. M., Berliocchi, L., Piomelli, D., and Tassorelli, C. (2017). Inhibition of monoacylglycerol lipase: another signalling pathway for potential therapeutic targets in migraine? *Cephalalgia*. doi: 10.1177/0333102417727537. [Epub ahead of print].
- Greco, R., Gasperi, V., Maccarrone, M., and Tassorelli, C. (2010a). The endocannabinoid system and migraine. *Exp. Neurol.* 224, 85–91. doi: 10.1016/j.expneurol.2010.03.029
- Greco, R., Gasperi, V., Sandrini, G., Bagetta, G., Nappi, G., Maccarrone, M., et al. (2010b). Alterations of the endocannabinoid system in an animal model of migraine: evaluation in cerebral areas of rat. *Cephalalgia* 30, 296–302. doi: 10.1111/j.1468-2982.2009.01924.x
- Greco, R., Mangione, A. S., Sandrini, G., Maccarrone, M., Nappi, G., and Tassorelli, C. (2011a). Effects of anandamide in migraine: data from an animal model. *J. Headache Pain* 12, 177–183. doi: 10.1007/s10194-010-0274-4
- Greco, R., Mangione, A. S., Sandrini, G., Nappi, G., and Tassorelli, C. (2014). Activation of CB2 receptors as a potential therapeutic target for migraine: evaluation in an animal model. *J. Headache Pain* 15:14. doi: 10.1186/1129-2377-15-14
- Greco, R., Meazza, C., Mangione, A. S., Allena, M., Bolla, M., Amantea, D., et al. (2011b). Temporal profile of vascular changes induced by systemic nitroglycerin in the meningeal and cortical districts. *Cephalalgia* 31, 190–198. doi: 10.1177/0333102410379887
- Gulyas, A. I., Cravatt, B. F., Bracey, M. H., Dinh, T. P., Piomelli, D., Boscia, F., et al. (2004). Segregation of two endocannabinoid-hydrolyzing enzymes into pre- and postsynaptic compartments in the rat hippocampus, cerebellum and amygdala. *Eur. J. Neurosci.* 20, 441–458. doi: 10.1111/j.1460-9568.2004.03428.x
- Hua, T., Vemuri, K., Pu, M., Qu, L., Han, G. W., Wu, Y., et al. (2016). Crystal structure of the human Cannabinoid Receptor CB1. *Cell* 167, 750.e14–762.e14. doi: 10.1016/j.cell.2016.10.004
- Jayamanne, A., Greenwood, R., Mitchell, V. A., Aslan, S., Piomelli, D., and Vaughan, C. W. (2006). Actions of the FAAH inhibitor URB597 in neuropathic and inflammatory chronic pain models. *Br. J. Pharmacol.* 147, 281–288. doi: 10.1038/sj.bjp.0706510
- Jhaveri, M. D., Sagar, D. R., Elmes, S. J., Kendall, D. A., and Chapman, V. (2007). Cannabinoid CB2 receptor-mediated anti-nociception in models of acute and chronic pain. *Mol. Neurobiol.* 36, 26–35. doi: 10.1007/s12035-007-8007-7
- Jin, X. H., Okamoto, Y., Morishita, J., Tsuboi, K., Tonai, T., and Ueda, N. (2007). Discovery and characterization of a Ca²⁺-independent phosphatidylethanolamine N-acyltransferase generating the anandamide precursor and its congeners. *J. Biol. Chem.* 282, 3614–3623. doi: 10.1074/jbc.M606369200
- Juhász, G., Csepány, E., Magyar, M., Edes, A. E., Eszlari, N., Hullam, G., et al. (2017). Variants in the CNR1 gene predispose to headache with nausea in the presence of life stress. *Genes Brain Behav.* 16, 384–393. doi: 10.1111/gbb.12352
- Kano, M., Ohno-Shosaku, T., Hashimoto, Y., Uchigashima, M., and Watanabe, M. (2009). Endocannabinoid-mediated control of synaptic transmission. *Physiol. Rev.* 89, 309–380. doi: 10.1152/physrev.00019.2008
- Katona, I., and Freund, T. F. (2008). Endocannabinoid signaling as a synaptic circuit breaker in neurological disease. *Nat. Med.* 14, 923–930. doi: 10.1038/nm.f.1869
- Kazemi, H., Rahgozar, M., Speckmann, E. J., and Gorji, A. (2012). Effect of cannabinoid receptor activation on spreading depression. *Iran. J. Basic Med. Sci.* 15, 926–936.
- Kelly, S., and Chapman, V. (2001). Selective cannabinoid CB1 receptor activation inhibits spinal nociceptive transmission *in vivo*. *J. Neurophysiol.* 86, 3061–3064. doi: 10.1152/jn.2001.86.6.3061
- Kendall, D. A., and Yudowski, G. A. (2017). Cannabinoid receptors in the central nervous system: their signaling and roles in disease. *Front. Cell. Neurosci.* 10:294. doi: 10.3389/fncel.2016.00294
- Khurana, L., Mackie, K., Piomelli, D., and Kendall, D. A. (2017). Modulation of CB1 cannabinoid receptor by allosteric ligands: pharmacology and therapeutic opportunities. *Neuropharmacology* 124, 3–12. doi: 10.1016/j.neuropharm.2017.05.018
- Kinsey, S. G., Long, J. Z., Cravatt, B. F., and Lichtman, A. H. (2010). Fatty acid amide hydrolase and monoacylglycerol lipase inhibitors produce anti-allodynic effects in mice through distinct cannabinoid receptor mechanisms. *J. Pain* 11, 1420–1428. doi: 10.1016/j.jpain.2010.04.001
- Kinsey, S. G., Long, J. Z., O'Neal, S. T., Abdullah, R. A., Poklis, J. L., Boger, D. L., et al. (2009). Blockade of endocannabinoid-degrading enzymes attenuates neuropathic pain. *J. Pharmacol. Exp. Ther.* 330, 902–910. doi: 10.1124/jpet.109.155465
- Knight, Y. E., Classey, J. D., Lasalandra, M. P., Akerman, S., Kowacs, F., Hoskin, K. L., et al. (2005). Patterns of fos expression in the rostral medulla and caudal pons evoked by noxious craniovascular stimulation and periaqueductal gray stimulation in the cat. *Brain Res.* 1045, 1–11. doi: 10.1016/j.brainres.2005.01.091
- La Rana, G., Russo, R., D'Agostino, G., Sasso, O., Raso, G. M., Iacono, A., et al. (2008). AM404, an anandamide transport inhibitor, reduces plasma extravasation in a model of neuropathic pain in rat: role for cannabinoid receptors. *Neuropharmacology* 54, 521–529. doi: 10.1016/j.neuropharm.2007.10.021
- Liu, J., Wang, L., Harvey-White, J., Osei-Hyiaman, D., Razdan, R., Gong, Q., et al. (2006). A biosynthetic pathway for anandamide. *Proc. Natl. Acad. Sci. U.S.A.* 103, 13345–13350. doi: 10.1073/pnas.0601832103
- Lodola, A., Castelli, R., Mor, M., and Rivara, S. (2015). Fatty acid amide hydrolase inhibitors: a patent review (2009–2014). *Expert Opin. Ther. Pat.* 25, 1247–1266. doi: 10.1517/13543776.2015.1067683
- Manzanares, J., Corchero, J., Romero, J., Fernandez-Ruiz, J. J., Ramos, J. A., and Fuentes, J. A. (1998). Chronic administration of cannabinoids regulates proenkephalin mRNA levels in selected regions of the rat brain. *Brain Res. Mol. Brain Res.* 55, 126–132. doi: 10.1016/S0169-328X(97)00371-9
- McKinney, M. K., and Cravatt, B. F. (2005). Structure and function of fatty acid amide hydrolase. *Annu. Rev. Biochem.* 74, 411–432. doi: 10.1146/annurev.biochem.74.082803.133450
- Mendiguren, A., Aostri, E., and Pineda, J. (2018). Regulation of noradrenergic and serotonergic systems by cannabinoids: relevance to cannabinoid-induced effects. *Life Sci.* 192, 115–127. doi: 10.1016/j.lfs.2017.11.029
- Messlinger, K., Suzuki, A., Pawlak, M., Zehnter, A., Schmidt, R. F., and Sánchez-Blázquez, P. (2000). Involvement of nitric oxide in the modulation of

- dural arterial blood flow in the rat. *Br. J. Pharmacol.* 129, 1397–1404. doi: 10.1038/sj.bjp.0703220
- Miquel, M. C., Emerit, M. B., Nosjean, A., Simon, A., Rumajogee, P., Brisorgueil, M. J., et al. (2002). Differential subcellular localization of the 5-HT₃-As receptor subunit in the rat central nervous system. *Eur. J. Neurosci.* 15, 449–457. doi: 10.1046/j.0953-816x.2001.01872.x
- Moldrich, G., and Wenger, T. (2000). Localization of the CB1 cannabinoid receptor in the rat brain. An immunohistochemical study. *Peptides* 21, 1735–1742. doi: 10.1016/S0196-9781(00)00324-7
- Moskowitz, M. A. (1993). Neurogenic inflammation in the pathophysiology and treatment of migraine. *Neurology* 43, S16–S20.
- Nackley, A. G., Suplita, R. L., and Hohmann, A. G. (2003). A peripheral cannabinoid mechanism suppresses spinal fos protein expression and pain behavior in a rat model of inflammation. *Neuroscience* 117, 659–670. doi: 10.1016/S0306-4522(02)00870-9
- Nackley, A. G., Zvonok, A. M., Makriyannis, A., and Hohmann, A. G. (2004). Activation of cannabinoid CB2 receptors suppresses C-fiber responses and windup in spinal wide dynamic range neurons in the absence and presence of inflammation. *J. Neurophysiol.* 92, 3562–3574. doi: 10.1152/jn.00886.2003
- Nagy-Grócz, G., Tar, L., Bohár, Z., Fejes-Szabó, A., Laborc, K. F., Spekker, E., et al. (2016). The modulatory effect of anandamide on nitroglycerin-induced sensitization in the trigeminal system of the rat. *Cephalalgia* 36, 849–861. doi: 10.1177/0333102415613766
- Nasirinezhad, F., Jergova, S., Pearson, J. P., and Sagen, J. (2015). Attenuation of persistent pain-related behavior by fatty acid amide hydrolase (FAAH) inhibitors in a rat model of HIV sensory neuropathy. *Neuropharmacology* 95, 100–109. doi: 10.1016/j.neuropharm.2014.11.024
- Nozaki, C., Markert, A., and Zimmer, A. (2015). Inhibition of FAAH reduces nitroglycerin-induced migraine-like pain and trigeminal neuronal hyperactivity in mice. *Eur. Neuropsychopharmacol.* 25, 1388–1396. doi: 10.1016/j.euroneuro.2015.04.001
- Ohooha, D. C., Knable, M. B., Wolf, S. S., Kleinman, J. E., and Hyde, T. M. (1994). The subnuclear distribution of 5-HT₃ receptors in the human nucleus of the solitary tract and other structures of the caudal medulla. *Brain Res.* 637, 222–226. doi: 10.1016/0006-8993(94)91236-X
- Okamoto, Y., Wang, J., Morishita, J., and Ueda, N. (2007). Biosynthetic pathways of the endocannabinoid anandamide. *Chem. Biodivers.* 4, 1842–1857. doi: 10.1002/cbdv.200790155
- Ong, J. J. Y., and De Felice, M. (2017). Migraine treatment: current acute medications and their potential mechanisms of action. *Neurotherapeutics*. doi: 10.1007/s13311-017-0592-1. [Epub ahead of print].
- Park, J. W., Shin, H. E., Kim, J. S., and Lee, K. S. (2008). Assessing migraine disability by diary-based measurement: relationship to the characteristics of individual headache attacks. *Eur. J. Neurol.* 15, 817–821. doi: 10.1111/j.1468-1331.2008.02204.x
- Parker, L. A., Rock, E. M., and Limebeer, C. L. (2011). Regulation of nausea and vomiting by cannabinoids. *Br. J. Pharmacol.* 163, 1411–1422. doi: 10.1111/j.1476-5381.2010.01176.x
- Patricelli, M. P., and Cravatt, B. F. (1999). Fatty acid amide hydrolase competitively degrades bioactive amides and esters through a nonconventional catalytic mechanism. *Biochemistry* 38, 14125–14130. doi: 10.1021/bi991876p
- Perrotta, A., Arce-Leal, N., Tassorelli, C., Gasperi, V., Sances, G., Blandini, F., et al. (2012). Acute reduction of anandamide-hydrolase (FAAH) activity is coupled with a reduction of nociceptive pathways facilitation in medication-overuse headache subjects after withdrawal treatment. *Headache* 52, 1350–1361. doi: 10.1111/j.1526-4610.2012.02170.x
- Pertwee, R. G. (2001). Cannabinoid receptors and pain. *Prog. Neurobiol.* 63, 569–611. doi: 10.1016/S0301-0082(00)00031-9
- Piomelli, D., and Sasso, O. (2014). Peripheral gating of pain signals by endogenous lipid mediators. *Nat. Neurosci.* 17, 164–174. doi: 10.1038/nn.3612
- Piomelli, D., Tarzia, G., Duranti, A., Tontini, A., Mor, M., Compton, T. R., et al. (2006). Pharmacological profile of the selective FAAH inhibitor KDS-4103 (URB597). *CNS Drug Rev.* 12, 21–38. doi: 10.1111/j.1527-3458.2006.00021.x
- Puente, N., Cui, Y., Lassalle, O., Lafourcade, M., Georges, F., Venance, L., et al. (2011). Polymodal activation of the endocannabinoid system in the extended amygdala. *Nat. Neurosci.* 14, 1542–1547. doi: 10.1038/nn.2974
- Quartillo, A., Mata, H. P., Ibrahim, M. M., Vanderah, T. W., Porreca, F., Makriyannis, A., et al. (2003). Inhibition of inflammatory hyperalgesia by activation of peripheral CB2 cannabinoid receptors. *Anesthesiology* 99, 955–960. doi: 10.1097/0000542-200310000-00031
- Ramachandran, R., Bhatt, D. K., Ploug, K. B., Hay-Schmidt, A., Jansen-Olesen, I., Gupta, S., et al. (2014). Nitric oxide synthase, calcitonin gene-related peptide and NK-1 receptor mechanisms are involved in GTN-induced neuronal activation. *Cephalalgia* 34, 136–147. doi: 10.1177/0333102413502735
- Reuter, U., Bolay, H., Jansen-Olesen, I., Chiarugi, A., Sanchez del Rio, M., Letourneau, R., et al. (2001). Delayed inflammation in rat meninges: implications for migraine pathophysiology. *Brain* 124, 2490–2502. doi: 10.1093/brain/124.12.2490
- Rhyne, D. N., Anderson, S. L., Gedde, M., and Borgelt, L. M. (2016). Effects of medical Marijuana on migraine headache frequency in an adult population. *Pharmacotherapy* 36, 505–510. doi: 10.1002/phar.1673
- Rossi, C., Pini, L. A., Cupini, M. L., Calabresi, P., and Sarchielli, P. (2008). Endocannabinoids in platelets of chronic migraine patients and medication-overuse headache patients: relation with serotonin levels. *Eur. J. Clin. Pharmacol.* 64, 1–8. doi: 10.1007/s00228-007-0391-4
- Sarchielli, P., Pini, L. A., Coppola, F., Rossi, C., Baldi, A., Mancini, M. L., et al. (2007). Endocannabinoids in chronic migraine: CSF findings suggest a system failure. *Neuropsychopharmacology* 32, 1384–1390. doi: 10.1038/sj.npp.1301246
- Shao, Z., Yin, J., Chapman, K., Grzemska, M., Clark, L., Wang, J., et al. (2016). High-resolution crystal structure of the human CB1 cannabinoid receptor. *Nature*. doi: 10.1038/nature20613. [Epub ahead of print].
- Slivicki, R. A., Xu, Z., Kulkarni, P. M., Pertwee, R. G., Mackie, K., Thakur, G. A., et al. (2017). Positive allosteric modulation of cannabinoid receptor type 1 suppresses pathological pain without producing tolerance or dependence. *Biol. Psychiatry*. doi: 10.1016/j.biopsych.2017.06.032. [Epub ahead of print].
- Urquhart, P., Nicolau, A., and Woodward, D. F. (2015). Endocannabinoids and their oxygenation by cyclo-oxygenases, lipoxygenases and other oxygenases. *Biochim. Biophys. Acta* 1851, 366–376. doi: 10.1016/j.bbali.2014.12.015
- Van der Schueren, B. J., Van Laere, K., Gérard, N., Bormans, G., and De Hoon, J. N. (2012). Intercalated type 1 cannabinoid receptor binding is increased in female migraine patients. *Headache* 52, 433–440. doi: 10.1111/j.1526-4610.2011.02030.x
- Van Sickle, M. D., Oland, L. D., Ho, W., Hillard, C. J., Mackie, K., Davison, J. S., et al. (2001). Cannabinoids inhibit emesis through CB1 receptors in the brainstem of the ferret. *Gastroenterology* 121, 767–774. doi: 10.1053/gast.2001.28466
- Volfe, Z., Dvilansky, A., and Nathan, I. (1985). Cannabinoids block release of serotonin from platelets induced by plasma from migraine patients. *Int. J. Clin. Pharmacol. Res.* 5, 243–246.
- Wilkerson, J. L., Ghosh, S., Mustafa, M., Abdullah, R. A., Niphakis, M. J., and Cabrera, R. (2017). The endocannabinoid hydrolysis inhibitor SA-57: intrinsic antinociceptive effects, augmented morphine-induced antinociception, and attenuated heroin seeking behavior in mice. *Neuropharmacology* 114, 156–167. doi: 10.1016/j.neuropharm.2016.11.015
- Zubrzycki, M., Janecka, A., Liebold, A., Ziegler, M., and Zubrzycka, M. (2017). Effects of centrally administered endocannabinoids and opioids on orofacial pain perception in rats. *Br. J. Pharmacol.* 174, 3780–3789. doi: 10.1111/bph.13970

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Greco, Demartini, Zanaboni, Piomelli and Tassorelli. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.