



Role of Dendritic Cells in Natural Immune Control of HIV-1 Infection

Enrique Martin-Gayo^{1*} and Xu G. Yu^{2*}

¹ Hospital Universitario de la Princesa, Universidad Autónoma de Madrid, Madrid, Spain, ² Ragon Institute of MGH, MIT, and Harvard, Massachusetts General Hospital, Harvard Medical School, Boston, MA, United States

OPEN ACCESS

Edited by:

Paul Urquhart Cameron,
The University of Melbourne, Australia

Reviewed by:

Phillippe Benaroch,
Centre National de la Recherche
Scientifique (CNRS), France
Laura Fantuzzi,
Istituto Superiore di Sanità (ISS), Italy

*Correspondence:

Enrique Martin-Gayo
enrique.martin@uam.es
Xu G. Yu
xyu@mgh.harvard.edu

Specialty section:

This article was submitted to
Viral Immunology,
a section of the journal
Frontiers in Immunology

Received: 21 March 2019

Accepted: 22 May 2019

Published: 06 June 2019

Citation:

Martin-Gayo E and Yu XG (2019) Role
of Dendritic Cells in Natural Immune
Control of HIV-1 Infection.
Front. Immunol. 10:1306.
doi: 10.3389/fimmu.2019.01306

Dendritic cells (DCs) are professional antigen-presenting cells that link innate and adaptive immunity and are critical for the induction of protective immune responses against pathogens. Proportions of these cells are markedly decreased in the blood of untreated HIV-1-infected individuals, suggesting they might be intrinsically involved in HIV-1 pathogenesis. However, despite several decades of active research, the precise role and contribution of these cells to protective or detrimental host responses against HIV-1 are still remarkably unclear. Recent studies have shown that DCs possess a fine-tuned machinery to recognize HIV-1 replication products through a variety of innate pathogen sensing mechanisms, which may be instrumental for generating both cellular and humoral protective immune responses in persons who naturally control HIV-1 replication. Yet, dysregulated and abnormal activation of DCs might also contribute to sustained inflammation and immune activation accelerating disease progression during chronic progressive infection. Emerging data also suggest that DCs can influence the induction of potent broadly-neutralizing antibodies, and may, for this reason, have to be considered as important components of future HIV-1 vaccination strategies. Apart from their involvement in antiviral host immunity, at least a subgroup of DCs seem intrinsically susceptible to HIV-1 infection and may serve as a viral target cell population. Indeed recent studies suggest that specific DC subpopulations residing in the genital mucosa are preferentially infected by HIV-1 and play an active role in sexual transmission; therefore, DCs may contribute to viral dissemination and possible persistence of the viral reservoirs through either direct or indirect mechanisms. Here, we analyze the distinct and partially opposing roles of DCs during HIV-1 disease pathogenesis, with a focus on implications of DC biology natural immune control and HIV cure research efforts.

Keywords: dendritic cell, HIV-1 controller, IFN, Tfh, bNAbs, vaccine

INTRODUCTION

Dendritic cells (DCs) represent a heterogeneous family of immune cells that link innate and adaptive immunity. The main function of these innate cells is to capture, process, and present antigens to adaptive immune cells and mediate their polarization into effector cells (1). DCs can be subdivided in two main subtypes: plasmacytoid (pDC) and myeloid (mDC) DCs, which specialize in the recognition of different pathogen associated molecular patterns (PAMPs) due to the unique distribution of Pattern Recognition Receptors (PRR), such as toll-like receptors, C-type lectins and intracellular nucleic acid sensors (2–4). As a result, mDCs and pDCs can efficiently induce CD4⁺ and CD8⁺ T cell responses against different types of pathogens. In addition, both mDCs




FIGURE 1 | Schematic representation of factors in human mDC and pDC contributing to immune control vs. progression of HIV-1 infection.

recent works in the SIV model have suggested that additional cell types might be responsible for abnormal activation of type I IFN responses at later stages of progressive infection (29). Together, cumulative information from recent studies suggests that DC distribution and function might be critically altered during HIV-1 infection and that preservation of physiological DC distribution and function is associated with immune control of the infection.

DCs AS VEHICLES FOR HIV-1 TRANSMISSION AND DISSEMINATION

During the last few years, several studies have shown that DCs have the ability to transfer HIV-1 particles to target CD4⁺ T cells and facilitate their infection, in a process known as trans-infection (30). This phenomenon starts with the transference of HIV-1 virions to pockets in the membrane of DCs, where they accumulate and are subsequently actively transferred to T cells through virological synapses (31). The stability of such transmission events depends on the expression of adhesion molecules, such as Intercellular Adhesion Molecule (ICAM) (32) and the actin assembly machinery (33). In order to transfer viruses to CD4⁺ T cells, DCs require the capture of HIV-1 particles through the lectin Dendritic Cell Intercellular Adhesion Molecule-3-Grabbing Non-integrin (DC-SIGN) (34, 35), and the Sialic acid-binding Immunoglobulin-type Lectin 1 (SIGLEC-1) receptor (36). Recent reports have shown that the ability of DCs to facilitate HIV-1 trans-infection is acquired upon activation with inflammatory molecules associated with poor HIV-1 prognosis, such as IFNα and LPS (37). These

stimuli have been shown to induce SIGLEC-1 expression and therefore, enhance the capture and transfer of viral particles. Notably, circulating mDCs but not pDCs facilitate trans-infection of HIV-1 (32). Therefore, increased basal levels of immune activation and high-level viremia might contribute to disease progression through facilitation of viral trans-infection by mDCs. Consistent with a role of mDCs in supporting HIV-1 trans-infection in lymphoid tissues, it was shown that depletion of lymph node-resident mDCs in tissue-suspension cultures reduced the efficiency of HIV-1 infection of CD4⁺ T cells (38). However, mutations of SIGLEC-1, which naturally occur in small proportions of individuals, did not seem to provide protection from HIV-1 infection or attenuation of HIV-1 disease progression (39), suggesting that classical, SIGLEC-1-independent HIV-1 dissemination within in the host remains the predominant mechanisms fueling viral infection *in vivo*.

DC SUSCEPTIBILITY TO HIV-1 INFECTION AND HOST RESTRICTION FACTORS

Most DCs express the coreceptor CD4 (40), and therefore are in principle susceptible to infection with HIV-1. However, DCs seem to represent a more hostile and restrictive environment for HIV-1 than CD4⁺ T cells, for reasons that are not completely clear. While initial studies suggested that monocyte derived DCs (MDDCs) are highly resistant to infection with HIV-1 (41), primary mDCs are able to support some levels of HIV-1 replication, at least *in vitro* (42–44).

The main restriction factor that limits HIV-1 replication in MDDCs and mDCs seems to be the cytoplasmic protein SAM domain and HD domain-containing protein 1 (SAMHD1), which is highly expressed in myeloid cells and is able to block HIV-1 replication at the retro-transcription level by depleting endogenous intracellular pools of dNTPs (45), and by directly degrading viral RNA (46). While it is clear that SAMHD1 is a key factor limiting replication of HIV-1 in MDDCs (47) and inhibiting further spread of virions to T cells (48), recent studies demonstrated that MDDCs can actually support productive infection with HIV-1 to a certain degree, despite high levels of expression of this restriction factor (49). The functional ability of SAMHD1 to restrict HIV-1 replication is regulated by phosphorylation mediated by host kinases from the cyclin-dependent kinase family (50). Interestingly, the functionally active, de-phosphorylated form of SAMHD1 is preferentially found in primary DCs isolated *ex vivo* from human blood, which potentially could contribute to a higher resistance to infection (51). However, it is unclear whether restriction of HIV-1 by SAMHD1 in mDCs might truly benefit the host, since restriction of HIV-1 replication via SAMHD1 may impair cytoplasmic viral immune recognition in mDCs and impair their ability to prime HIV-1-specific T cells. On the other hand, interactions of mDCs with T cells induce downregulation of SAMHD1 expression (52), allowing human primary mDCs to be more permissive to infection (44). Importantly, recent data indicate that primary CD1c⁺ and CD141⁺ mDC subtypes might differ in their susceptibility to HIV-1 infection. In this regard, expression of the endosomal protein RAB15 prevents fusion of viral particles in CD141⁺ mDCs and induces a higher level of cell-intrinsic resistance to infection with HIV-1 and HIV-2 compared to CD1c⁺ mDCs (53). Further proof for the susceptibility of primary mDCs to HIV-1 infection was provided by a recent study identifying a distinct population of CD1a⁺ mDCs residing in the vaginal mucosa, which supported CCR5-tropic but not CXCR4-tropic HIV-1 replication, in contrast to vaginal Langerhans cells (LC). These data suggest that these vaginal mDCs might play an active role in the selection of transmitted viral variants during heterosexual HIV-1 acquisition (54).

In the context of immune control of HIV-1 infection, recent studies suggest that monocytes and mDCs from HIV-1 controllers restrict early HIV-1 replication steps, specifically at the level of viral integration (44, 55, 56) while restriction of viral reverse transcription is less obvious, possibly due to lower induction of SAMHD1 expression in HIV-1 controllers upon exposure to HIV-1. This specific replicative pattern of HIV-1 may enable enhanced cytoplasmic sensing of accumulated HIV-1 reverse transcripts, which represent the primary substrate for innate immune recognition, and facilitate antigen processing and presentation (44, 56). Interestingly, although SAMHD1 is thought to be an interferon inducible gene, DCs and CD4⁺ T cells fail to induce its expression in the presence of type I IFNs (57). Therefore, higher permissiveness of mDCs from controllers to viral reverse transcription may represent a key element for supporting cytoplasmic detection of HIV-1 and for inducing potent cell-intrinsic responses that lead to the effective activation of HIV-1-specific T cells (Figure 1).

Although SAMHD1 is recognized as a critical host factor limiting HIV-1 replication in myeloid cells, alternative SAMHD1-independent restriction mechanisms might also be playing a role in effective immunological control of HIV-1 replication. Among them, recognition of the HIV-1 capsid by cyclophilin A (41, 58) and TRIM5 α (59, 60) or endogenous levels of β -catenin (52), could be actively contributing to block HIV-1 replication in myeloid cells. In addition, some studies suggest that HIV-1 could trigger TLR activation in DCs (61). Indeed, activation of MDDCs through TLR4 and TLR3 resulted in inhibition of HIV-1 replication steps in DC, while simultaneously increasing their ability to prime HIV-specific CD8⁺ T cells (62). Therefore, TLR-dependent activation of DC could play a relevant role for inducing highly-functional cellular immune responses against HIV-1. Supporting this idea, polymorphisms in the TLR3 gene confer resistance to HIV-1 infection (63). In fact, it was recently suggested that TLR activation could be playing an active role in the detection of HIV-1 by primary CD141⁺ mDCs (53). Therefore, more studies are required to investigate the mutual interplay between viral restriction in DCs and immune control of HIV-1, and to determine the contribution of myeloid cells to persisting viral reservoirs during suppressive antiretroviral therapy.

INNATE IMMUNE RESPONSES TO HIV-1 IN DCs

DCs are, in principle, capable of inducing secretion of type I IFNs upon recognition of viral nucleic acids, which subsequently leads to transcription of interferon stimulated genes (ISGs) and the upregulation of class II HLA and costimulatory molecules. As a result of such cell-intrinsic, IFN-dependent immune responses, mature DCs become more restrictive for viral replication, while the expression of molecules involved in antigen presentation and co-stimulation is increased. Whether mDCs can induce secretion of type I IFNs in response to HIV-1 is still highly controversial. In MDDCs, HIV-1 seems to be able to induce expression of several IFN-related genes in the absence of actual production of IFN α/β due to the selective activation of IRF-1 mediated signaling instead of inducing phosphorylation of IRF3, which is known to be required for induction of type I IFNs (64). However, the intracellular DNA sensor cGAS is expressed by myeloid cells (65) and is able of producing cGAMP second messengers upon recognition of HIV-1 DNA (66), leading to the activation of the sensor STING and the signal transducer TBK-1, which promote IFN β production (67, 68). Thus, primary DCs are, in principle, able to sense and induce type I IFN upon exposure to cytoplasmic HIV-1 DNA. In fact, activation of cGAS seems to be required for the transcription of IFN β by primary mDCs and MDDCs in the context of HIV-1 and other viral infections (44, 69, 70). This DNA-dependent mechanism of viral sensing leading to type I IFN responses might be more active in human CD1c⁺ mDCs compared to CD141⁺ mDCs (53). Interestingly, cGAS triggers TLR9-independent activation of primary pDCs in response to intracellular DNA (71, 72). However, current phenotypic

markers for pDC identify a heterogeneous cell population that, in addition to *bona fide* pDC, contains pre-DC precursors of mDCs, which could also be differentially contributing to the observed responses to HIV-1 (73). Such heterogeneity could be the result of different pre-pDC and/or pDCs originated from either lymphoid or myeloid precursors with different functional properties (74–76). Therefore, more studies are required to elucidate the contribution of TLR-independent sensing of HIV-1 in pDCs. Finally, activation of the cGAS pathway by HIV-1 might involve interactions with additional host factors such as the newly identified NONO protein, which apparently is able to bind cGAS and the HIV-1 capsid and facilitate innate sensing of HIV-1 DNA in dendritic cells (77).

Importantly, preserved or enhanced induction of IFN responses has been described in both primary pDCs (23, 24) and mDCs (44) from HIV-1 controllers exposed to HIV-1. A recent single-cell RNAseq study has identified a highly functional population of CD64^{Hi}CD86^{Hi} PD-L1^{Hi} mDCs characterized by a strong type I IFN signature that is induced more efficiently in HIV-1 controllers than in progressors or healthy individuals in response to HIV-1 (78). The induction of such highly functional mDCs depended on the activation of TBK-1, which acts downstream of several intracellular sensing pathways including cGAS and TLR-3. Therefore, enhanced innate recognition of HIV-1 by both pDCs and mDCs might be a contributing factor to develop effective HIV-1 specific immunity in these individuals (Figure 1). However, HIV-1 might have evolved to minimize such mechanisms of viral DNA recognition, since HIV-1 Vif and Vpr are capable of inactivating TBK-1 which is downstream of the cGAS-STING pathway (79). Therefore, additional mechanisms such as alterations in the activation threshold of intracellular sensing pathways might be playing a role in DCs from controllers. In addition to sensing of viral DNA by cGAS, viral immune recognition in DCs could be connected with the RIG-I pathway, which may also contribute to activation of DCs in response to HIV-1 (80). In fact, communication and collaboration between RIG-I and DNA sensing pathways has been reported to amplify innate immune responses against intracellular viral DNA (81, 82). Although no genetic alterations in genes encoding for innate immune sensors for HIV-1 have been found in GWAS studies including large HIV-1-infected populations, a more targeted analysis of innate immune genes may in the future allow to identify immunogenetic polymorphisms in the innate immune system that facilitate innate immune sensing and natural viral control in specific subgroups of HIV-1 controllers.

ANTIGEN PRESENTING CELL FUNCTION OF DCs AND ADAPTIVE IMMUNITY AGAINST HIV-1

Given associations between the polyfunctionality of T cell responses and natural progression of HIV-1 infection (8), several studies have focused on the function of DCs as professional antigen presenting cells (APC) and how these cells are involved in the priming of adaptive immune cells. As mentioned before,

both mDCs and pDCs can respond and mature to a certain degree in response to HIV-1, but may become exhausted and hyporesponsive during chronic progressive infection, which might impact their antigen-presenting cell function (83). In the case of pDCs, infection with HIV-1 seems to turn these cells more tolerogenic, and increase their potential to drive polarization of CD4⁺ T cells into immunosuppressive T regulatory cells (84). On the other hand, although pDCs can activate CD8⁺ T cells through cross-presentation (85), no studies have yet analyzed the impact of pDCs on the priming of HIV-1-specific cytotoxic CD8⁺ T cell responses. In contrast, while circulating mDCs from healthy individuals are functionally incapable of efficiently priming T cells *in vitro* after exposure to HIV-1 (86), effective antigen presenting functions of mDCs from HIV-1 elite controllers is associated with enhanced abilities to prime HIV-specific CD8⁺ T cells in these patients (44, 78). In addition to mDCs, recent *in vitro* studies have shown that MDDCs can acquire HIV-1 antigens from Langerhans cells, become activated and induce cross-presentation to CD8⁺ T cells (87), suggesting that these cells may also be potentially able to prime protective HIV-1-specific cytotoxic CD8⁺ T cells. In addition, independent studies have shown that MDDCs can mediate cross-presentation of immuno-dominant HIV-1 peptides and activate HIV-1-specific CD8⁺ T cells (88). However, MDDC are not a physiological DC subset present and in fact more closely resemble inflammatory DCs (89). Nevertheless, a recent evaluation suggested that primary CD141⁺ mDCs, obtain HIV-1 antigens from infected CD1c⁺ mDCs for cross-presentation to CD8⁺ T cells (53). Interestingly, DCs infected with HIV-1 can also present endogenous viral peptides and mediate activation of HIV-1-specific CD4⁺ T cells (90). Therefore, mDCs and MDDCs might be involved in the priming of effective HIV-1-specific T cell responses observed in controllers.

Although highly functional HIV-1-specific CD8⁺ T cell responses were identified as the main correlate of antiviral immune defense (91), the discovery of broadly neutralizing antibodies (bNAbs) against multiple strains of HIV-1 (92, 93), has led to a great interest in understanding their potential contribution to spontaneous immunological control of HIV-1. Recent works have indeed identified a subpopulation of HIV-1 viremic controllers who develop bNAbs in the absence of high levels of viremia or immune activation (94). In previous studies in viremic HIV-1-infected progressors, the induction of bNAbs was associated with the presence of CXCR5⁺ PD-1⁺ T follicular helper cells (Tfh) in the blood (95). Although Tfh cells facilitate B cell maturation and immunoglobulin class switching in lymphoid tissue (96), peripheral CXCR5⁺ PD-1⁺ CD4⁺ T lymphocytes have been proposed to act as peripheral counterparts of Tfh cells (pTfh) (95, 97) and could serve as a peripheral biomarker of high germinal center Tfh cell activity. Therefore, the priming of Tfh cells by mDC might be important in HIV-1 controllers capable of inducing antibodies with broader neutralizing activity. Supporting this possibility, mDCs from controller neutralizers are more efficient in priming CD4⁺ T cells into long lived PD-1^{Lo} Tfh precursors, which can differentiate into functional PD-1^{Hi} Tfh effector cells upon antigenic stimulation (98). Importantly, higher frequencies of PD-1^{Lo} Tfh precursors in the blood are

correlated with higher breadth of Ab neutralization in this subset of controllers. Compatible with an indirect role of DCs for influencing humoral immunity through polarization of Tfh, mDCs from controller neutralizers are characterized by high levels of CD40, a molecule previously involved in Tfh cell differentiation (99), and display distinct transcriptional patterns that differ from those present in CD64^{Hi} PD-L1^{Hi} mDCs from elite controllers with high CD8⁺ T cell priming potential (98). Therefore, these findings might suggest a range of functional specializations of mDCs from controllers that may contribute to immune viral control through different immune mechanisms. To which degree individual DC subpopulations influence other components of the innate and adaptive immune system and contribute to control of HIV-1 is still an open question that requires further study.

CONCLUSIONS

In this review, we have summarized recent advances in understanding DC biology in the context of HIV-1 immune control. While studies have revealed multiple mechanisms by which DCs might contribute to controlling HIV-1 (Figure 1), future studies will be necessary to evaluate the complexity of individual DC subsets in promoting beneficial versus detrimental

effects during HIV-1 infection. Similarly, our knowledge about the intrinsic ability of pDCs and mDCs to sense and respond to HIV-1 has greatly improved over the last few years, but translating this insight into improved and more specific adjuvants for future preventive and therapeutic HIV-1 vaccines represents a considerable challenge. The development to new humanized animal models that recapitulate human DC biology will likely be critical to identify effective vaccination strategies based on DCs. Together, DCs are emerging as critical players of effective immune responses in HIV-1 and a closer understanding of these cells might contribute to the development of novel effective vaccines or immunotherapies.

AUTHOR CONTRIBUTIONS

EM-G and XY conceived and wrote the manuscript.

FUNDING

XY is supported by NIH Grants AI078799, HL134539, AI116228, AI125109, DA047034, AI117841, AI114235, AI135940. EM-G was supported by the Programa Atracción de Talento Modalidad 1 from Comunidad de Madrid (2017-T1/BMD-5396) and the NIH R21grant program (AI140930).

REFERENCES

- Shortman K, Liu YJ. Mouse and human dendritic cell subtypes. *Nat Rev Immunol.* (2002) 2:151–61. doi: 10.1038/nri746
- Jarrossay D, Napolitani G, Colonna M, Sallusto F, Lanzavecchia A. Specialization and complementarity in microbial molecule recognition by human myeloid and plasmacytoid dendritic cells. *Eur J Immunol.* (2001) 31:3388–93. doi: 10.1002/1521-4141(200111)31:11<3388::AID-IMMU3388>3.0.CO;2-Q
- Szabo A, Rajnavolgyi E. Collaboration of Toll-like and RIG-I-like receptors in human dendritic cells: tRIGgering antiviral innate immune responses. *Am J Clin Exp Immunol.* (2013) 2:195–207.
- Unterholzner L. The interferon response to intracellular DNA: why so many receptors? *Immunobiology.* (2013) 218:1312–21. doi: 10.1016/j.imbio.2013.07.007
- Marcenaro E, Carlomagno S, Pesce S, Moretta A, Sivori S. NK/DC crosstalk in anti-viral response. *Adv Exp Med Biol.* (2012) 946:295–308. doi: 10.1007/978-1-4614-0106-3_17
- Blankson JN. Effector mechanisms in HIV-1 infected elite controllers: highly active immune responses? *Antiviral Research.* (2010) 85:295–302. doi: 10.1016/j.antiviral.2009.08.007
- Hersperger AR, Martin JN, Shin LY, Sheth PM, Kovacs CM, Cosma GL, et al. Increased HIV-specific CD8⁺ T-cell cytotoxic potential in HIV elite controllers is associated with T-bet expression. *Blood.* (2011) 117:3799–808. doi: 10.1182/blood-2010-12-322727
- Saez-Cirion A, Lacabaratz C, Lambotte O, Versmisse P, Urrutia A, Boufassa F, et al. HIV controllers exhibit potent CD8 T cell capacity to suppress HIV infection *ex vivo* and peculiar cytotoxic T lymphocyte activation phenotype. *Proc Natl Acad Sci USA.* (2007) 104:6776–81. doi: 10.1073/pnas.0611244104
- Kadowaki N, Ho S, Antonenko S, Malefyt RW, Kastelein RA, Bazan F, et al. Subsets of human dendritic cell precursors express different toll-like receptors and respond to different microbial antigens. *J Exp Med.* (2001) 194:863–9. doi: 10.1084/jem.194.6.863
- Tavano B, Galao RP, Graham DR, Neil SJ, Aquino VN, Fuchs D, et al. Ig-like transcript 7, but not bone marrow stromal cell antigen 2 (also known as HM1.24, tetherin, or CD317), modulates plasmacytoid dendritic cell function in primary human blood leukocytes. *J Immunol.* (2013) 190:2622–30. doi: 10.4049/jimmunol.1202391
- Barron MA, Blyveis N, Palmer BE, MaWhinney S, Wilson CC. Influence of plasma viremia on defects in number and immunophenotype of blood dendritic cell subsets in human immunodeficiency virus 1-infected individuals. *J Infect Dis.* (2003) 187:26–37. doi: 10.1086/345957
- Dillon SM, Robertson KB, Pan SC, Mawhinney S, Meditz AL, Folkvord JM, et al. Plasmacytoid and myeloid dendritic cells with a partial activation phenotype accumulate in lymphoid tissue during asymptomatic chronic HIV-1 infection. *J Acq Immune Defic Syndromes.* (2008) 48:1–12. doi: 10.1097/QAI.0b013e3181664b60
- Sabado RL, O'Brien M, Subedi A, Qin L, Hu N, Taylor E, et al. Evidence of dysregulation of dendritic cells in primary HIV infection. *Blood.* (2010) 116:3839–52. doi: 10.1182/blood-2010-03-273763
- Diao Y, Geng W, Fan X, Cui H, Sun H, Jiang Y, et al. Low CD1c + myeloid dendritic cell counts correlated with a high risk of rapid disease progression during early HIV-1 infection. *BMC Infect Dis.* (2015) 15:342. doi: 10.1186/s12879-015-1092-8
- Malleret B, Karlsson I, Maneglier B, Brochard P, Delache B, Andrieu T, et al. Effect of SIVmac infection on plasmacytoid and CD1c+ myeloid dendritic cells in cynomolgus macaques. *Immunology.* (2008) 124:223–33. doi: 10.1111/j.1365-2567.2007.02758.x
- Donaghy H, Pozniak A, Gazzard B, Qazi N, Gilmour J, Gotch F, et al. Loss of blood CD11c(+) myeloid and CD11c(-) plasmacytoid dendritic cells in patients with HIV-1 infection correlates with HIV-1 RNA virus load. *Blood.* (2001) 98:2574–6. doi: 10.1182/blood.V98.8.2574
- Lehmann C, Jung N, Forster K, Koch N, Leifeld L, Fischer J, et al. Longitudinal analysis of distribution and function of plasmacytoid dendritic cells in peripheral blood and gut mucosa of HIV infected patients. *J Infect Dis.* (2014) 209:940–9. doi: 10.1093/infdis/jit612
- Cunningham CR, Champhekar A, Tullius MV, Dillon BJ, Zhen A, de la Fuente JR, et al. Type I and type II interferon coordinately regulate suppressive dendritic cell fate and function during viral persistence. *PLoS Pathogens.* (2016) 12:e1005356. doi: 10.1371/journal.ppat.1005356
- Carranza P, Del Rio Estrada PM, Diaz Rivera D, Ablanedo-Terrazas Y, Reyes-Teran G. Lymph nodes from HIV-infected individuals harbor mature

- dendritic cells and increased numbers of PD-L1+ conventional dendritic cells. *Human Immunol.* (2016) 77:584–93. doi: 10.1016/j.humimm.2016.05.019
20. Huang J, Burke P, Yang Y, Seiss K, Beamon J, Cung T, et al. Soluble HLA-G inhibits myeloid dendritic cell function in HIV-1 infection by interacting with leukocyte immunoglobulin-like receptor B2. *J Virol.* (2010) 84:10784–91. doi: 10.1128/JVI.01292-10
 21. Huang J, Burke PS, Cung TD, Pereyra F, Toth I, Walker BD, et al. Leukocyte immunoglobulin-like receptors maintain unique antigen-presenting properties of circulating myeloid dendritic cells in HIV-1-infected elite controllers. *J Virol.* (2010) 84:9463–71. doi: 10.1128/JVI.01009-10
 22. Lehmann C, Harper JM, Taubert D, Hartmann P, Fatkenheuer G, Jung N, et al. Increased interferon alpha expression in circulating plasmacytoid dendritic cells of HIV-1-infected patients. *J Acquired Immune Defic Syndr.* (2008) 48:522–30. doi: 10.1097/QAI.0b013e31817f97cf
 23. Barblu L, Machmach K, Gras C, Delfraissy JF, Boufassa F, Leal M, et al. Plasmacytoid dendritic cells (pDCs) from HIV controllers produce interferon-alpha and differentiate into functional killer pDCs under HIV activation. *J Infect Dis.* (2012) 206:790–801. doi: 10.1093/infdis/jis384
 24. Machmach K, Leal M, Gras C, Viciano P, Genebat M, Franco E, et al. Plasmacytoid dendritic cells reduce HIV production in elite controllers. *J Virol.* (2012) 86:4245–52. doi: 10.1128/JVI.07114-11
 25. Saidi H, Bras M, Formaglio P, Melki MT, Charbit B, Herbeuval JP, et al. HMGB1 Is Involved in IFN-alpha production and TRAIL expression by HIV-1-exposed plasmacytoid dendritic cells: impact of the crosstalk with NK cells. *PLoS Pathog.* (2016) 12:e1005407. doi: 10.1371/journal.ppat.1005407
 26. Herbeuval JP, Grivel JC, Boasso A, Hardy AW, Choungnet C, Dolan MJ, et al. CD4+ T-cell death induced by infectious and noninfectious HIV-1: role of type 1 interferon-dependent, TRAIL/DR5-mediated apoptosis. *Blood.* (2005) 106:3524–31. doi: 10.1182/blood-2005-03-1243
 27. Lichtner M, Rossi R, Rizza MC, Mengoni F, Sauzullo I, Massetti AP, et al. Plasmacytoid dendritic cells count in antiretroviral-treated patients is predictive of HIV load control independent of CD4+ T-cell count. *Curr HIV Res.* (2008) 6:19–27. doi: 10.2174/157016208783571937
 28. Royle CM, Graham DR, Sharma S, Fuchs D, Boasso A. HIV-1 and HIV-2 differentially mature plasmacytoid dendritic cells into IFN-producing cells or APCs. *J Immunol.* (2014) 193:3538–48. doi: 10.4049/jimmunol.1400860
 29. Kader M, Smith AP, Guiducci C, Wonderlich ER, Normolle D, Watkins SC, et al. Blocking TLR7- and TLR9-mediated IFN-alpha production by plasmacytoid dendritic cells does not diminish immune activation in early SIV infection. *PLoS Pathog.* (2013) 9:e1003530. doi: 10.1371/journal.ppat.1003530
 30. Kijewski SD, Gummuluru S. A mechanistic overview of dendritic cell-mediated HIV-1 trans infection: the story so far. *Future Virol.* (2015) 10:257–69. doi: 10.2217/fvl.15.2
 31. Dale BM, Alvarez RA, Chen BK. Mechanisms of enhanced HIV spread through T-cell virological synapses. *Immunol Rev.* (2013) 251:113–24. doi: 10.1111/imr.12022
 32. Groot F, van Capel TM, Kapsenberg ML, Berkhout B, de Jong EC. Opposing roles of blood myeloid and plasmacytoid dendritic cells in HIV-1 infection of T cells: transmission facilitation versus replication inhibition. *Blood.* (2006) 108:1957–64. doi: 10.1182/blood-2006-03-010918
 33. Menager MM, Littman DR. Actin dynamics regulates dendritic cell-mediated transfer of HIV-1 to T cells. *Cell.* (2016) 164:695–709. doi: 10.1016/j.cell.2015.12.036
 34. Geijtenbeek TB, Kwon DS, Torensma R, van Vliet SJ, van Duinhoven GC, Middel J, et al. DC-SIGN, a dendritic cell-specific HIV-1-binding protein that enhances trans-infection of T cells. *Cell.* (2000) 100:587–97. doi: 10.1016/S0092-8674(00)80694-7
 35. Kwon DS, Gregorio G, Bitton N, Hendrickson WA, Littman DR. DC-SIGN-mediated internalization of HIV is required for trans-enhancement of T cell infection. *Immunity.* (2002) 16:135–44. doi: 10.1016/S1074-7613(02)00259-5
 36. Izquierdo-Useros N, Lorizate M, McLaren PJ, Telenti A, Krausslich HG, Martinez-Picado J. HIV-1 capture and transmission by dendritic cells: the role of viral glycolipids and the cellular receptor Siglec-1. *PLoS Pathog.* (2014) 10:e1004146. doi: 10.1371/journal.ppat.1004146
 37. Pino M, Erkizia I, Benet S, Erikson E, Fernandez-Figueroas MT, Guerrero D, et al. HIV-1 immune activation induces Siglec-1 expression and enhances viral trans-infection in blood and tissue myeloid cells. *Retrovirology.* (2015) 12:37. doi: 10.1186/s12977-015-0160-x
 38. Reyes-Rodriguez AL, Reuter MA, McDonald D. Dendritic cells enhance HIV infection of memory CD4(+) T cells in human lymphoid tissues. *AIDS Res Human Retroviruses.* (2016) 32:203–10. doi: 10.1089/aid.2015.0235
 39. Martinez-Picado J, McLaren PJ, Erkizia I, Martin MP, Benet S, Rotger M, et al. Identification of Siglec-1 null individuals infected with HIV-1. *Nat Comm.* (2016) 7:12412. doi: 10.1038/ncomms12412
 40. Patterson S, Rae A, Hockey N, Gilmour J, Gotch F. Plasmacytoid dendritic cells are highly susceptible to human immunodeficiency virus type 1 infection and release infectious virus. *J Virol.* (2001) 75:6710–3. doi: 10.1128/JVI.75.14.6710-6713.2001
 41. Manel N, Hogstad B, Wang Y, Levy DE, Unutmaz D, Littman DR. A cryptic sensor for HIV-1 activates antiviral innate immunity in dendritic cells. *Nature.* (2010) 467:214–7. doi: 10.1038/nature09337
 42. Smed-Sorensen A, Lore K, Vasudevan J, Louder MK, Andersson J, Mascola JR, et al. Differential susceptibility to human immunodeficiency virus type 1 infection of myeloid and plasmacytoid dendritic cells. *J Virol.* (2005) 79:8861–9. doi: 10.1128/JVI.79.14.8861-8869.2005
 43. Cameron PU, Handley AJ, Baylis DC, Solomon AE, Bernard N, Purcell DF, et al. Preferential infection of dendritic cells during human immunodeficiency virus type 1 infection of blood leukocytes. *J Virol.* (2007) 81:2297–306. doi: 10.1128/JVI.01795-06
 44. Martin-Gayo E, Buzon MJ, Ouyang Z, Hickman T, Cronin J, Pimenova D, et al. Potent cell-intrinsic immune responses in dendritic cells facilitate HIV-1-specific T cell immunity in HIV-1 elite controllers. *PLoS Pathog.* (2015) 11:e1004930. doi: 10.1371/journal.ppat.1004930
 45. Lahouassa H, Daddacha W, Hofmann H, Ayinde D, Logue EC, Dragin L, et al. SAMHD1 restricts the replication of human immunodeficiency virus type 1 by depleting the intracellular pool of deoxynucleoside triphosphates. *Nat Immunol.* (2012) 13:223–8. doi: 10.1038/ni.2236
 46. Ryoo J, Choi J, Oh C, Kim S, Seo M, Kim SY, et al. The ribonuclease activity of SAMHD1 is required for HIV-1 restriction. *Nat Med.* (2014) 20:936–41. doi: 10.1038/nm.3626
 47. Hrecka K, Hao C, Gierszewska M, Swanson SK, Kesik-Brodacka M, Srivastava S, et al. Vpx relieves inhibition of HIV-1 infection of macrophages mediated by the SAMHD1 protein. *Nature.* (2011) 474:658–61. doi: 10.1038/nature10195
 48. Puigdomenech I, Casartelli N, Porrot F, Schwartz O. SAMHD1 restricts HIV-1 cell-to-cell transmission and limits immune detection in monocyte-derived dendritic cells. *J Virol.* (2013) 87:2846–56. doi: 10.1128/JVI.02514-12
 49. Hertoghs N, van der Aar AM, Setiawan LC, Kootstra NA, Gringhuis SI, Geijtenbeek TB. SAMHD1 degradation enhances active suppression of dendritic cell maturation by HIV-1. *J Immunol.* (2015) 194:4431–7. doi: 10.4049/jimmunol.1403016
 50. Cribier A, Descours B, Valadao AL, Laguette N, Benkirane M. Phosphorylation of SAMHD1 by cyclin A2/CDK1 regulates its restriction activity toward HIV-1. *Cell Rep.* (2013) 3:1036–43. doi: 10.1016/j.celrep.2013.03.017
 51. Bloch N, O'Brien M, Norton TD, Polsky SB, Bhardwaj N, Landau NR. HIV type 1 infection of plasmacytoid and myeloid dendritic cells is restricted by high levels of SAMHD1 and cannot be counteracted by Vpx. *AIDS Res Human Retroviruses.* (2014) 30:195–203. doi: 10.1089/aid.2013.0119
 52. Aljawai Y, Richards MH, Seaton MS, Narasipura SD, Al-Harhi L. beta-Catenin/TCF-4 signaling regulates susceptibility of macrophages and resistance of monocytes to HIV-1 productive infection. *Curr HIV Res.* (2014) 12:164–73. doi: 10.2174/1570162X12666140526122249
 53. Silvin A, Yu CI, Lahaye X, Imperatore F, Braut JB, Cardinaud S, et al. Constitutive resistance to viral infection in human CD141(+) dendritic cells. *Sci Immunol.* (2017) 2:eai8071. doi: 10.1126/sciimmunol.aai8071
 54. Pena-Cruz V, Agosto LM, Akiyama H, Olson A, Moreau Y, Larriex JR, et al. HIV-1 replicates and persists in vaginal epithelial dendritic cells. *J Clin Invest.* (2018) 128:3439–44. doi: 10.1172/JCI98943
 55. Saez-Cirion A, Hamimi C, Bergamaschi A, David A, Versmisse P, Melard A, et al. Restriction of HIV-1 replication in macrophages and CD4+ T cells from HIV controllers. *Blood.* (2011) 118:955–64. doi: 10.1182/blood-2010-12-327106
 56. Hamimi C, David A, Versmisse P, Weiss L, Bruel T, Zucman D, et al. Dendritic cells from HIV controllers have low susceptibility to HIV-1 infection *in vitro* but high capacity to capture HIV-1 particles. *PLoS ONE.* (2016) 11:e0160251. doi: 10.1371/journal.pone.0160251

57. St. Gelais C, de Silva S, Amie SM, Coleman CM, Hoy H, Hollenbaugh JA, et al. SAMHD1 restricts HIV-1 infection in dendritic cells (DCs) by dNTP depletion, but its expression in DCs and primary CD4+ T-lymphocytes cannot be upregulated by interferons. *Retrovirology*. (2012) 9:105. doi: 10.1186/1742-4690-9-105
58. Lahaye X, Satoh T, Gentili M, Cerboni S, Silvin A, Conrad C, et al. Nuclear envelope protein SUN2 promotes cyclophilin-a-dependent steps of HIV replication. *Cell Rep*. (2016) 15:879–92. doi: 10.1016/j.celrep.2016.03.074
59. Pertel T, Hausmann S, Morger D, Zuger S, Guerra J, Lascano J, et al. TRIM5 is an innate immune sensor for the retrovirus capsid lattice. *Nature*. (2011) 472:361–5. doi: 10.1038/nature09976
60. Portilho DM, Fernandez J, Ringeard M, Machado AK, Boulay A, Mayer M, et al. Endogenous TRIM5alpha function is regulated by SUMOylation and nuclear sequestration for efficient innate sensing in dendritic cells. *Cell Rep*. (2016) 14:355–69. doi: 10.1016/j.celrep.2015.12.039
61. Ben Haij N, Planes R, Leghmar K, Serrero M, Delobel P, Izopet J, et al. HIV-1 tat protein induces production of proinflammatory cytokines by human dendritic cells and monocytes/macrophages through engagement of TLR4-MD2-CD14 complex and activation of NF-kappaB pathway. *PLoS ONE*. (2015) 10:e0129425. doi: 10.1371/journal.pone.0129425
62. Cardinaud S, Urrutia A, Rouers A, Coulon PG, Kerveyan J, Richetta C, et al. Triggering of TLR-3,–4, NOD2, and DC-SIGN reduces viral replication and increases T-cell activation capacity of HIV-infected human dendritic cells. *Eur J Immunol*. (2017) 47:818–29. doi: 10.1002/eji.201646603
63. Sironi M, Biasin M, Cagliani R, Forni D, De Luca M, Saule I, et al. A common polymorphism in TLR3 confers natural resistance to HIV-1 infection. *J Immunol*. (2012) 188:818–23. doi: 10.4049/jimmunol.1102179
64. Harman AN, Lai J, Turville S, Samarajiwa S, Gray L, Marsden V, et al. HIV infection of dendritic cells subverts the IFN induction pathway via IRF-1 and inhibits type 1 IFN production. *Blood*. (2011) 118:298–308. doi: 10.1182/blood-2010-07-297721
65. Sun L, Wu J, Du F, Chen X, Chen ZJ. Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science*. (2013) 339:786–91. doi: 10.1126/science.1232458
66. Gao D, Wu J, Wu YT, Du F, Aroh C, Yan N, et al. Cyclic GMP-AMP synthase is an innate immune sensor of HIV and other retroviruses. *Science*. (2013) 341:903–6. doi: 10.1126/science.1240933
67. Zhang H, Tang K, Zhang Y, Ma R, Ma J, Li Y, et al. Cell-free tumor microparticle vaccines stimulate dendritic cells via cGAS/STING signaling. *Cancer Immunol Res*. (2015) 3:196–205. doi: 10.1158/2326-6066.CIR-14-0177
68. Shu C, Li X, Li P. The mechanism of double-stranded DNA sensing through the cGAS-STING pathway. *Cytokine Growth Factor Rev*. (2014) 25:641–8. doi: 10.1016/j.cytogfr.2014.06.006
69. Lahaye X, Satoh T, Gentili M, Cerboni S, Conrad C, Hurbain I, et al. The capsids of HIV-1 and HIV-2 determine immune detection of the viral cDNA by the innate sensor cGAS in dendritic cells. *Immunity*. (2013) 39:1132–42. doi: 10.1016/j.immuni.2013.11.002
70. Yoh SM, Schneider M, Seifried J, Soonthornvacharin S, Akleh RE, Olivieri KC, et al. PQBP1 is a proximal sensor of the cGAS-dependent innate response to HIV-1. *Cell*. (2015) 161:1293–305. doi: 10.1016/j.cell.2015.04.050
71. Bode C, Fox M, Tewary P, Steinhagen A, Ellerkmann RK, Klinman D, et al. Human plasmacytoid dendritic cells elicit a Type I Interferon response by sensing DNA via the cGAS-STING signaling pathway. *Eur J Immunol*. (2016) 46:1615–21. doi: 10.1002/eji.201546113
72. Paijo J, Doring M, Spanier J, Grabski E, Nooruzzaman M, Schmidt T, et al. cGAS senses human cytomegalovirus and induces type I interferon responses in human monocyte-derived cells. *PLoS Pathog*. (2016) 12:e1005546. doi: 10.1371/journal.ppat.1005546
73. See P, Dutertre CA, Chen J, Gunther P, McGovern N, Irac SE, et al. Mapping the human DC lineage through the integration of high-dimensional techniques. *Science*. (2017) 356:eaag3009. doi: 10.1126/science.aag3009
74. Rodrigues PF, Alberti-Servera L, Eremin A, Grajales-Reyes GE, Ivanek R, Tussiwand R. Distinct progenitor lineages contribute to the heterogeneity of plasmacytoid dendritic cells. *Nat Immunol*. (2018) 19:711–22. doi: 10.1038/s41590-018-0136-9
75. Herman JS, Sagar, Grun D. FateID infers cell fate bias in multipotent progenitors from single-cell RNA-seq data. *Nat Methods*. (2018) 15:379–86. doi: 10.1038/nmeth.4662
76. Martin-Gayo E, Gonzalez-Garcia S, Garcia-Leon MJ, Murcia-Ceballos A, Alcaín J, Garcia-Peydro M, et al. Spatially restricted JAG1-Notch signaling in human thymus provides suitable DC developmental niches. *J Exp Med*. (2017) 214:3361–79. doi: 10.1084/jem.20161564
77. Lahaye X, Gentili M, Silvin A, Conrad C, Picard L, Jouve M, et al. NONO Detects the nuclear HIV capsid to promote cgas-mediated innate immune activation. *Cell*. (2018) 175:488–501.e22. doi: 10.1016/j.cell.2018.08.062
78. Martin-Gayo E, Cole MB, Kolb KE, Ouyang Z, Cronin J, Kazer SW, et al. A reproducibility-based computational framework identifies an inducible, enhanced antiviral state in dendritic cells from HIV-1 elite controllers. *Genome Biol*. (2018) 19:10. doi: 10.1186/s13059-017-1385-x
79. Harman AN, Nasr N, Feetham A, Galoyan A, Alshehri AA, Rambukwelle D, et al. HIV blocks interferon induction in human dendritic cells and macrophages by dysregulation of TBK1. *J Virol*. (2015) 89:6575–84. doi: 10.1128/JVI.00889-15
80. Berg RK, Melchjorsen J, Rintahaka J, Diget E, Soby S, Horan KA, et al. Genomic HIV RNA induces innate immune responses through RIG-I-dependent sensing of secondary-structured RNA. *PLoS ONE*. (2012) 7:e29291. doi: 10.1371/journal.pone.0029291
81. Ablasser A, Bauernfeind F, Hartmann G, Latz E, Fitzgerald KA, Hornung V. RIG-I-dependent sensing of poly(dA:dT) through the induction of an RNA polymerase III-transcribed RNA intermediate. *Nat Immunol*. (2009) 10:1065–72. doi: 10.1038/ni.1779
82. Ahlers LR, Bastos RG, Hiroyasu A, Goodman AG. Invertebrate iridescent virus 6, a DNA virus, stimulates a mammalian innate immune response through RIG-I-like receptors. *PLoS ONE*. (2016) 11:e0166088. doi: 10.1371/journal.pone.0166088
83. Kaushik S, Teque F, Patel M, Fujimura SH, Schmidt B, Levy JA. Plasmacytoid dendritic cell number and responses to Toll-like receptor 7 and 9 agonists vary in HIV Type 1-infected individuals in relation to clinical state. *AIDS Res Human Retroviruses*. (2013) 29:501–10. doi: 10.1089/aid.2012.0200
84. Manches O, Fernandez MV, Plumas J, Chaperot L, Bhardwaj N. Activation of the noncanonical NF-kappaB pathway by HIV controls a dendritic cell immunoregulatory phenotype. *Proc Natl Acad Sci USA*. (2012) 109:14122–7. doi: 10.1073/pnas.1204032109
85. Tel J, Schreiber G, Sittig SP, Mathan TS, Buschow SI, Cruz LJ, et al. Human plasmacytoid dendritic cells efficiently cross-present exogenous Ags to CD8+ T cells despite lower Ag uptake than myeloid dendritic cell subsets. *Blood*. (2013) 121:459–67. doi: 10.1182/blood-2012-06-435644
86. Granelli-Piperno A, Shimeliovich I, Pack M, Trumpfheller C, Steinman RM. HIV-1 selectively infects a subset of nonmaturing BDCA1-positive dendritic cells in human blood. *J Immunol*. (2006) 176:991–8. doi: 10.4049/jimmunol.176.2.991
87. van den Berg LM, Cardinaud S, van der Aar AM, Sprokholz JK, de Jong MA, Zijlstra-Willems EM, et al. Langerhans cell-dendritic cell cross-talk via langerin and hyaluronic acid mediates antigen transfer and cross-presentation of HIV-1. *J Immunol*. (2015) 195:1763–73. doi: 10.4049/jimmunol.1402356
88. Dinter J, Duong E, Lai NY, Berberich MJ, Kourjian G, Bracho-Sanchez E, et al. Variable processing and cross-presentation of HIV by dendritic cells and macrophages shapes CTL immunodominance and immune escape. *PLoS Pathog*. (2015) 11:e1004725. doi: 10.1371/journal.ppat.1004725
89. Segura E, Touzot M, Bohineust A, Cappuccio A, Chiochia G, Hosmalin A, et al. Human inflammatory dendritic cells induce Th17 cell differentiation. *Immunity*. (2013) 38:336–48. doi: 10.1016/j.immuni.2012.10.018
90. Coulon PG, Richetta C, Rouers A, Blanchet FP, Urrutia A, Guerbois M, et al. HIV-infected dendritic cells present endogenous MHC class II-restricted antigens to HIV-specific CD4+ T cells. *J Immunol*. (2016) 197:517–32. doi: 10.4049/jimmunol.1600286
91. Walker BD, Yu XG. Unravelling the mechanisms of durable control of HIV-1. *Nat Rev Immunol*. (2013) 13:487–98. doi: 10.1038/nri3478
92. Kwong PD, Mascola JR. Human antibodies that neutralize HIV-1: identification, structures, and B cell ontogenies. *Immunity*. (2012) 37:412–25. doi: 10.1016/j.immuni.2012.08.012
93. Burton DR, Mascola JR. Antibody responses to envelope glycoproteins in HIV-1 infection. *Nat Immunol*. (2015) 16:571–6. doi: 10.1038/ni.3158
94. Doria-Rose NA, Klein RM, Daniels MG, O'Dell S, Nason M, Lapedes A, et al. Breadth of human immunodeficiency virus-specific neutralizing activity in

- sera: clustering analysis and association with clinical variables. *J Virol.* (2010) 84:1631–6. doi: 10.1128/JVI.01482-09
95. Locci M, Havenar-Daughton C, Landais E, Wu J, Kroenke MA, Arlehamn CL, et al. Human circulating PD-1+CXCR3-CXCR5+ memory T_H cells are highly functional and correlate with broadly neutralizing HIV antibody responses. *Immunity.* (2013) 39:758–69. doi: 10.1016/j.immuni.2013.08.031
96. Breitfeld D, Ohl L, Kremmer E, Ellwart J, Sallusto F, Lipp M, et al. Follicular B helper T cells express CXC chemokine receptor 5, localize to B cell follicles, and support immunoglobulin production. *J Exp Med.* (2000) 192:1545–52. doi: 10.1084/jem.192.11.1545
97. Morita R, Schmitt N, Bentebibel SE, Ranganathan R, Bourdery L, Zurawski G, et al. Human blood CXCR5(+)CD4(+) T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion. *Immunity.* (2011) 34:108–21. doi: 10.1016/j.immuni.2011.01.009
98. Martin-Gayo E, Cronin J, Hickman T, Ouyang Z, Lindqvist M, Kolb KE, et al. Circulating CXCR5+CXCR3+PD-1^{lo} T_H-like cells in HIV-1 controllers with neutralizing antibody breadth. *JCI Insight.* (2017) 2:e89574. doi: 10.1172/jci.insight.89574
99. Ma CS, Suryani S, Avery DT, Chan A, Nanan R, Santner-Nanan B, et al. Early commitment of naive human CD4(+) T cells to the T follicular helper (T_{FH}) cell lineage is induced by IL-12. *Immunol Cell Biol.* (2009) 87:590–600. doi: 10.1038/icc.2009.64

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Martin-Gayo and Yu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.